

Management of multi-drug resistant *Helicobacter pylori* infection by supplementary, complementary and alternative medicine; a review

Nasim Rezaeimanesh¹, Nastaran Farzi², Samira Pirmanesh³, Saeed Emami⁴, Abbas Yadegar²

¹Students Research Committee, Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Foodborne and Waterborne Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ABSTRACT

Helicobacter pylori is recognized as the most common bacterial pathogens colonizing the gastric epithelium of nearly half of the world's population. This bacterium is the main etiological cause of gastroduodenal ulcers, and more importantly as the substantial risk factor for development of gastric cancer. The emergence and rapid increase in the prevalence of multi-drug resistant phenotypes have posed major pitfalls in effectiveness of various treatment regimens and eradication strategies against *H. pylori* infections. Several natural products and supplementary food components have been reported to have established anti-*H. pylori* activity. Herein, we review the application and efficacy of some specific natural products and foodstuffs such as milk, bee products (honey and propolis), fish oil, vitamins C and E, and also a nickel free-diet used as anti-*H. pylori* alternative treatment regimens.

Keywords: *Helicobacter pylori*, Drug resistance, Dietary supplementation, Anti-inflammatory, Eradication therapy

(Please cite as: Rezaeimanesh N, Farzi N, Pirmanesh S, Emami S, Yadegar A. Management of multi-drug resistant *Helicobacter pylori* infection by supplementary, complementary and alternative medicine; a review. *Gastroenterol Hepatol Bed Bench* 2017; 10 (Suppl. 1):S8-S14).

Introduction

Helicobacter pylori is a spiral bacterium that colonize in the human gastric epithelium. The bacterium causes different precancerous lesions like gastritis, atrophy, intestinal metaplasia and dysplasia, and is the strongest known risk factor for gastric cancer (GC) (1-3). *H. pylori* infects more than half of the people globally, and the prevalence of *H. pylori* infection is highly variable across different countries; for example, high prevalence is observed in developing countries (~80%) in comparison to

developed countries with about 30-50% of the population (4). The bacterium usually is acquired in childhood and can persist for lifetime in the host stomach. *H. pylori* pathogenesis is mediated by a complex interplay between various bacterial virulence factors, host genetic predisposition, and environmental factors (5-7). *H. pylori* is also known as one of the most genetically diverse bacterial species that presents various virulence genotypes responsible for different gastric diseases (8-10). The emergence and more importantly the increasing prevalence of multi-drug resistant strains of *H. pylori* has led to reduced success in different treatment regimens (11-17). Several natural products and supplementary nutrients have been reported to have established antimicrobial activity against *H. pylori* infection (Table 1). Here, we tried to have a short

Received: 29 September 2017 Accepted: 18 November 2017
Reprint or Correspondence: Abbas Yadegar, PhD.
Foodborne and Waterborne Diseases Research Center,
Research Institute for Gastroenterology and Liver
Diseases, Shahid Beheshti University of Medical Sciences,
Tehran, Iran.
E-mail: babak_y1983@yahoo.com

overview on the application and efficacy of some natural products and supplementary compounds used as anti-*H. pylori* alternative treatments.

Dairy, bee products and fish oil,

Vitamins, Nickel free-diet

Milk

Milk, especially the human milk, has long been identified as one of the natural products encompassing high nutritional values as well as antimicrobial effects against a variety of infections. Before the discovery of antibiotics, a non-specific therapy named protein therapy, used the antibacterial properties of milk protein contents (18). Moreover, nutrient fortification of human milk with medium-chain triglycerides (MCT) and iron has been also applied as an acceptable supplement therapy for feeding preterm infants (19).

Lactoferrin is a multifunctional iron-binding glycoprotein with potent antibacterial and immunomodulatory properties against several bacterial pathogens (20, 21). It is released in the human colostrums with highest concentration in comparison with human milk and cow milk (20). It is also present in saliva, tears, seminal fluid and immune cells like neutrophils (20, 22). Lactoferrin has been shown to have inhibitory effects on growth of a number of bacterial pathogens including *Streptococcus mutans*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Neisseria meningitidis*, *Escherichia coli* and *H. pylori* (21, 23). The antimicrobial activity of lactoferrin is well studied and is likely attributed to its high affinity for iron, functioning as an iron chelator that sequesters iron elements from bacterial access (22, 24).

There are some in vivo studies showing that lactoferrin can improve *H. pylori* eradication rate in humans and mouse models (25-28). In another in vitro study, Akedo et al. showed anti-*H. pylori* property of cow's milk (29). Moreover, in a survey on 482 children aged 0-12-years-old, Okuda and et al. assessed the relation between breast feeding and *H. pylori* infection, and proposed breast feeding can be a natural way to protect children from infection due to anti-adhesive property of lactoferrin, sialyllactose and oligosaccharides present in breast milk (30). In 2013, a review study suggested that fermented milk-based probiotic preparations and bovine lactoferrin can be effective for *H. pylori* eradication (31).

Honey

In the traditional medicine, honey was used as an antimicrobial substance for treatment of infectious diseases, and also gastrointestinal disorders like dyspepsia, gastritis, peptic ulcer disease, and liver disease (27-29). Additionally, it is reported that honey can accelerate wound healing process and was used to dress wounds and burns (32). The antibacterial property of honey varies due to its color and floral sources. Red honeys have shown more potent antibacterial properties than white honeys, which this property depends on different phenolic acid contents of various kinds of honeys (33, 34). The bactericidal and bacteriostatic potency of honey is broad-spectrum, and is effective against several bacterial agents such as *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *E. coli* and *Streptococcus pyogenes* (34). It has been proposed that antibacterial activity of honey is mainly due to its high osmotic and acidity, hydrogen peroxide production, flavonoids and over-mentioned phenolic acids content (35-38).

Some studies revealed that high osmotic and high acidity effects of honey can inhibit the urease activity of *H. pylori* (37, 38). Abdel-Latif and colleagues investigated the molecular mechanisms by which natural honey may inhibit *H. pylori* infection. They reported that Manuka honey can inhibit *H. pylori* by suppression of *H. pylori* induced NF- κ B and AP-1 activation, and down regulation of COX-2 expression in gastric epithelial cells (35). Another in vivo study in Bulgaria showed the anti-*H. pylori* effect of honey via its high osmotic and acidity effect (36).

Propolis

Propolis is a resinous bee product that contains plant resin, bee enzymes and wax (39, 40). It is reported to be an antioxidant, antibacterial, antifungal, anti-inflammatory, antiproliferative and antidiabetic substance. The natural composition of propolis varies due to its original floral sources, and contains different amounts of phenolic compounds (39-41). The anti-*H. pylori* properties of propolis may be due to its phenolic substances such as flavonoids, phenolic acids and their esters like caffeic acid phenethyl ester (CAPE) and chrysin (40-42). Baltas and colleagues studied the anti-*H. pylori* effect of 15 ethanol extracts of propolis and reported that all extracts inhibited *H. pylori* J99 strain by urease inhibition (41). In 2013, Cui et al. assayed

S10 Management of multi-drug resistant *Helicobacter pylori* infection

different propolis phenolic compounds for *H. pylori* peptide deformylase (HpPDF) inhibition, which is necessary for *H. pylori* perpetuity and persistence (40).

Fish oil

From many years ago, essential oils were among the most useful components in traditional medicine around the world and their activity against *H. pylori* have been delineated (43, 44). Fish oil (Eicosapen) includes 33.5% omega-3-fatty acids with a variety of immunomodulating effects, which has bacteriostatic effect on *H. pylori* (45-49). Moreover, it has been reported that omega-3-fatty acids declined the secretion of gastric acid in healthy volunteers (50). It has been proposed that the inhibitory effects of fish oil on *H. pylori* may be due to: 1) direct inhibition or killing the bacteria (49), 2) inhibition of bacterial adhesion to gastric epithelium, and 3) inhibition of the *H. pylori*-induced inflammatory pathways (51, 52).

Vitamins

Vitamin C, an acidic molecule, is one of the most important component of living tissues. Two forms of vitamin C including: AA (ascorbic acid) and DHA (dehydroascorbic acid), the reduced and oxidized form, respectively, which can convert to each other. Inside the cell, DHA is immediately converted to AA in presence of glutathione or other thiols as electron donors via the specific enzyme systems like DHA reductase, glutaredoxins and protein disulfide isomerase (53, 54). Unfortunately, stability of AA and DHA are low and have a rapid wild irreversible hydrolysis particularly at a pH > 4 (55). Vitamins C and E have been studied to show their antioxidant effect for eradication of *H. pylori* infection (56, 57). It seems vitamins C and E break the microenvironment created by *H. pylori* or directly inhibit bacteria. Additionally, the detriment of antioxidants on colonization and proliferation of *H. pylori* have been shown (58-60).

In a study by Sezikli et al., they showed that under the oxidative stress vitamins C and E were effective in eradication of *H. pylori* infection (56). In another work, administration of high dose vitamin C treatment had inhibitory effects on *H. pylori* growth (61). In a study by Demirci and coworkers, the effect of vitamins C and E supplementation along with triple and quadruple eradication regimens was assessed using 400 *H. pylori* infected patients. They showed that *H. pylori* eradication rate was 56% for smokers and 94% for non-

smokers. The success rate of *H. pylori* eradication for smokers was lower than non-smokers (62). Zojaji et.al, also reported that addition of vitamin C to *H. pylori* treatment regimen including amoxicillin, metronidazole, and bismuth increased the eradication rate among the infected patients (63).

Nickel free-diet

Nickel is a metallic element that is widely found in almost all kinds of diets (64). It is abundant in fruits and vegetables like apricots, figs, pears, plums, raisins, pineapples, cabbage, onions, beans, lentils, potatoes, peas, tomatoes, spinach, cauliflower, asparagus, corn and margarine. This element is also present in different kinds of nuts including almonds, peanuts, walnuts, hazelnuts, and cocoa as well as some sea foods like lobster, mussels, oysters and plaice (65, 66). However, it seems that nickel is not essential for humans, but it is necessary for *H. pylori* colonization because of its important role in activation of *H. pylori* urease and hydrogenase enzymes. So, there is no competition between *H. pylori* and human body for nickel access (64). According to these facts, Campanale and colleagues in 2014 designed a pilot study to investigate the effect of a nickel free-diet on the eradication of *H. pylori* infection. In their in vivo study, 52 participants with *H. pylori* infection were divided into two groups: standard triple therapy and standard triple therapy with nickel free-diet. In the second group, the participants were prohibited to consume foods with high quantity of nickel for 4 weeks. They found that addition of nickel free-diet to standard triple therapy can significantly promote *H. pylori* eradication rate (64).

Discussion

Currently, the treatment of all symptomatic *H. pylori*-infected patients is less probable, and could rapidly increase the emergence and prevalence of multi-drug resistant strains in the community. Moreover, despite the availability of several therapeutic strategies for *H. pylori*-induced gastric diseases, the bacterial eradication is very challenging and none of the treatment regimens appear to be ideal. Therefore, the application of relatively low-cost natural products and foodstuffs with established anti-*H. pylori* activity seems to be promising as alternative medicine and adjuvant

Table 1. Specific food and natural products with potential anti-*H. pylori* activities.

Food types	Active components	Putative anti- <i>H. pylori</i> properties	Testing methods	Year/country	Ref.
	Lactoferrin, sialyllactose, oligosaccharides	Inhibition of <i>H. pylori</i> attachment	<i>H. pylori</i> stool antigen assay (HpSA)	2001/Japan	30
Milk	Lactoferrin adsorbed into biomimetic hydroxyapatite nanocrystals	Iron chelating and sequestration	Inhibition zone assay and the bacterial load were measured in orally <i>H. pylori</i> -infected BALB/c mice using SYBR Green I quantitative real-time PCR assay	Italy/2016	67
	Bovine milk glycoproteins and glycoconjugates and lactoferrin	Iron deprivation, decreasing gastric colonization of <i>H. pylori</i> and inflammation score	Growth inhibition assay, haemagglutination inhibition assay and adherence assay	Sweden/2001	28
	Hydrogen peroxide and phytochemicals (flavonoids and phenolic acids)	High osmotic effect, pH (high acidity)	¹³ C UBT	Bulgaria/2015	36
	Flavonoids and phenolic acids	Inhibition of bacterial urease activity	Urease activity assay by spectrophotometry	South Africa/2014	38
Honey		High osmotic effect, pH (high acidity)	Hole plate diffusion method and microbroth dilution method	Cameroon/2013	37
	Hydrogen peroxide and phytochemicals (flavonoids and phenolic acids)	Inhibition of <i>H. pylori</i> -induced NF-κB and AP-1 activation and downregulation of COX-2	Electrophoretic mobility shift assay (EMSA), cell viability assay and cytotoxicity assay	Egypt/2016	35
	Flavanoids, phenolic acids, vitamins, trace elements, amino acids, proteins, certain enzymes including glucose oxidase, invertase and catalase	expression, growth inhibition			
Propolis	Flavonoids, Phenolic compounds, Caffeic acid phenethyl ester and chrysin	Urease inhibition	Agar-well diffusion method and urease inhibition assay	Turkey/2016	41
	Phenolic compounds, Caffeic acid phenethyl ester	Inhibition of <i>H. pylori</i> peptide deformylase	Enzymatic activity of <i>HpPDF</i> was evaluated using a FDH coupled assay	China/2013	40
Fish oil	EPA, DHA, Omega 3 fatty acids	Direct inhibition of bacteria, antiadhesive activity, anti-inflammatory effect	Agar diffusion test (Kirby-Bauer method)	Italy/1999	52
Vitamin C	Ascorbic acid	<i>H. pylori</i> growth inhibition, urease inhibition, antioxidant effects	Urease test and histological examination (Giemsa staining)	Poland/1998	61
	Ascorbic acid, tocopherols and tocotrienols	Inhibition of <i>H. pylori</i> colonization, antioxidant effects	Rapid urease test, histopathological evaluation and UBT	Turkey/2015	62
Vitamins C and E	Ascorbic acid, tocopherols and tocotrienols	Antioxidant effects	Histologic examination, rapid urease test, ¹⁴ C-urea breath test, HpSA and Measurement of Total Antioxidant Capacity (TAC)	Turkey/2009	56

Abbreviations: UBT, urease breath test; *HpPDF*, *H. pylori* peptide deformylase; FDH, formate dehydrogenase; PCR, Polymerase chain reaction; PH, potential of hydrogen; NF-κB, nuclear factor kappa-light-chain-enhancer of activated B cells; AP-1, Activator protein 1; COX-2, Prostaglandin-endoperoxide synthase 2 (cyclooxygenase-2); EPA, Eicosapentaenoic acid; DHA, docosahexaenoic acid

S12 Management of multi-drug resistant Helicobacter pylori infection

therapy to manage the infections caused by antibiotic-resistant *H. pylori* strains. However, it is very important to evaluate the antibacterial effectiveness of different natural products and food components by both *in vitro* and *in vivo* experiments, especially in the clinical trials, to propose a potentially effective diet-based treatment regimen (68). Finally, further studies are needed to explore novel, local and natural therapeutics to be co-administrated with conventional antimicrobial agents as adjunctive therapy against *H. pylori* infections.

Acknowledgment

This study was financially supported by the Students Research Committee and Research & Technology Chancellor, and also by the Research Institute for Gastroenterology and Liver Diseases from Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Conflict of interests

The authors declare that they have no conflict of interest.

References

1. Shokrzadeh L, Baghaei K, Yamaoka Y, Dabiri H, Jafari F, Sahebkhietari N, et al. Analysis of 3'-end variable region of the *cagA* gene in *Helicobacter pylori* isolated from Iranian population. *J Gastroenterol Hepatol* 2010;25:172-77.
2. Baghaei K, Shokrzadeh L, Jafari F, Dabiri H, Yamaoka Y, Bolfion M, et al. Determination of *Helicobacter pylori* virulence by analysis of the *cag* pathogenicity island isolated from Iranian patients. *Dig Liver Dis* 2009;41:634-38.
3. Kao CY, Chen JW, Huang YT, Sheu SM, Sheu BS, Wu JJ. Genome sequence and annotation of *Helicobacter pylori* strain Hp238, isolated from a Taiwanese patient with mucosa-associated lymphoid tissue lymphoma. *Genome Announc* 2015;3:00006-15.
4. Sahebkhietari N, Nochi Z, Eslampour MA, Dabiri H, Bolfion M, Taherikalani M, et al. Characterization of *Staphylococcus aureus* strains isolated from raw milk of bovine subclinical mastitis in Tehran and Mashhad. *Acta Microbiol Immunol Hung* 2011;58:113-21.
5. Taremi M, Mehdi Soltan Dallal M, Gachkar L, MoezArdalan S, Zolfagharian K, Reza Zali M. Prevalence and antimicrobial resistance of *Campylobacter* isolated from retail raw chicken and beef meat, Tehran, Iran. *Int J Food Microbiol* 2006;108:401-403.
6. Yadegar A, Alebouyeh M, Lawson AJ, Mirzaei T, Nazemalhosseini Mojarad E, Zali MR. Differentiation of non-*pylori Helicobacter* species based on PCR-restriction

fragment length polymorphism of the 23S rRNA gene. *World J Microbiol Biotechnol* 2014;30:1909-17.

7. Yadegar A, Alebouyeh M, Zali MR. Analysis of the intactness of *Helicobacter pylori* *cag* pathogenicity island in Iranian strains by a new PCR-based strategy and its relationship with virulence genotypes and EPIYA motifs. *Infect Genet Evol* 2015;35:19-26.
8. Cover TL. *Helicobacter pylori* diversity and gastric cancer risk. *MBio* 2016;7:01869-15.
9. Gunaletchumy SP, Seevasant I, Tan MH, Croft LJ, Mitchell HM, Goh KL, et al. *Helicobacter pylori* genetic diversity and gastro-duodenal diseases in Malaysia. *Sci Rep* 2014;4:7431.
10. Farzi N, Malekian T, Alebouyeh M, Vaziri F, Zali MR. Genotype diversity and quasispecies development of *Helicobacter pylori* in a single host. *Jpn J Infect Dis* 2015;68:176-80.
11. Sugimoto M, Yamaoka Y. Virulence factor genotypes of *Helicobacter pylori* affect cure rates of eradication therapy. *Arch Immunol Ther Exp (Warsz)* 2009;57:45-56.
12. Boehnke KF, Valdivieso M, Bussalleu A, Sexton R, Thompson KC, Osorio S, et al. Antibiotic resistance among *Helicobacter pylori* clinical isolates in Lima, Peru. *Infect Drug Resist* 2017;10:85-90.
13. Megraud F. *H. pylori* antibiotic resistance: prevalence, importance, and advances in testing. *Gut* 2004;53:1374-84.
14. Vianna JS, Ramis IB, Ramos DF, VON GROLL A, Silva PEAd. Drug resistance in *Helicobacter pylori*. *Arq Gastroenterol* 2016;53:215-23.
15. Thung I, Aramin H, Vavinskaya V, Gupta S, Park J, Crowe S, et al. Review article: the global emergence of *Helicobacter pylori* antibiotic resistance. *Aliment Pharmacol Ther* 2016;43:514-33.
16. Shokrzadeh L, Alebouyeh M, Mirzaei T, Farzi N, Zali MR. Prevalence of multiple drug-resistant *Helicobacter pylori* strains among patients with different gastric disorders in Iran. *Microb Drug Resist* 2015;21:105-10.
17. Alebouyeh M, Yadegar A, Farzi N, Miri M, Zojaji H, Gharibi S, et al. Impacts of *H. pylori* mixed-infection and heteroresistance on clinical outcomes. *Gastroenterol Hepatol Bed Bench* 2015;8:S1-5.
18. Lahov E, Regelson W. Antibacterial and immunostimulating casein-derived substances from milk: casecidin, isracidin peptides. *Food Chem Toxicol* 1996;34:131-45.
19. Chan GM. Effects of powdered human milk fortifiers on the antibacterial actions of human milk. *J Perinatol* 2003;23:620-3.
20. Farnaud S, Evans RW. Lactoferrin--a multifunctional protein with antimicrobial properties. *Mol Immunol* 2003;40:395-405.
21. Jenssen H, Hancock RE. Antimicrobial properties of lactoferrin. *Biochimie* 2009;91:19-29.

22. Fahey JW, Stephenson KK, Wallace AJ. Dietary amelioration of *Helicobacter* infection. *Nutr Res* 2015;35:461-73.
23. Roşeanu A, Damian M, Evans RW. Mechanisms of the antibacterial activity of lactoferrin and lactoferrin-derived peptides. *Rom J Biochem* 2010;47:203-9.
24. Hołubiuk Ł, Imiela J. Diet and *Helicobacter pylori* infection. *Gastroenterol Rev* 2016;11:150-4.
25. Moosavi A, Haghighi A, Mojarad EN, Zayeri F, Alebouyeh M, Khazan H, et al. Genetic variability of *Blastocystis* sp. isolated from symptomatic and asymptomatic individuals in Iran. *Parasitol Res* 2012;111:2311-15.
26. Dial E, Romero J, Headon D, Lichtenberger L. Recombinant human lactoferrin is effective in the treatment of *Helicobacter felis*-infected mice. *J Pharm Pharmacol* 2000;52:1541-6.
27. Okuda M, Nakazawa T, Yamauchi K, Miyashiro E, Koizumi R, Booka M, et al. Bovine lactoferrin is effective to suppress *Helicobacter pylori* colonization in the human stomach: a randomized, double-blind, placebo-controlled study. *J Infect Chemother* 2005;11:265-9.
28. Wang X, Hirno S, Willen R, Wadström T. Inhibition of *Helicobacter pylori* infection by bovine milk glycoconjugates in a BALB/cA mouse model. *J Med Microbiol* 2001;50:430-5.
29. Akedo I, Tatsuta M, Narahara H, Iishi H, Uedo N, Yano H, et al. Prevention by bovine milk against *Helicobacter pylori*-associated atrophic gastritis through its adherence inhibition. *Hepatogastroenterology* 2003;51:277-81.
30. Okuda M, Miyashiro E, Koike M, Okuda S, Minami K, Yoshikawa N. Breast-feeding prevents *Helicobacter pylori* infection in early childhood. *Pediatr Int* 2001;43:714-5.
31. Sachdeva A, Rawat S, Nagpal J. Efficacy of fermented milk and whey proteins in *Helicobacter pylori* eradication: a review. *World J Gastroenterol* 2014;20:724-37.
32. Lusby P, Coombes A, Wilkinson J. Honey: a potent agent for wound healing? *J Wound Ostomy Continence Nurs* 2002;29:295-300.
33. Hegazi AG, Guthami FM, Gethami AF, Allah FM, Saleh AA, Fouad EA. Potential antibacterial activity of some Saudi Arabia honey. *Vet World* 2017;10:233-7.
34. Wasihun AG, Kasa BG. Evaluation of antibacterial activity of honey against multidrug resistant bacteria in Ayder Referral and Teaching Hospital, Northern Ethiopia. *SpringerPlus* 2016;5:842.
35. Abdel-Latif MM, Abouzied MM. Molecular mechanisms of natural honey against *H. pylori* infection via suppression of NF- κ B and AP-1 activation in gastric epithelial cells. *Arch Med Res* 2016;47:340-8.
36. Boyanova L, Ilieva J, Gergova G, Vladimirov B, Nikolov R, Mitov I. Honey and green/black tea consumption may reduce the risk of *Helicobacter pylori* infection. *Diagn Microbiol Infect Dis* 2015;82:85-6.
37. Manyi-Loh CE, Clarke AM, Green E, Ndip RN. Inhibitory and bactericidal activity of selected South African honeys and their solvent extracts against clinical isolates of *Helicobacter pylori*. *Pak J Pharm Sci* 2013;26:897-906.
38. Matongo F, Nwodo UU. In vitro assessment of *Helicobacter pylori* ureases inhibition by honey fractions. *Arch Med Res* 2014;45:540-6.
39. Boonsai P, Phuwapraisirisan P, Chanchao C. Antibacterial activity of a cardanol from Thai *Apis mellifera* propolis. *Int J Med Sci* 2014;11:327-36.
40. Cui K, Lu W, Zhu L, Shen X, Huang J. Caffeic acid phenethyl ester (CAPE), an active component of propolis, inhibits *Helicobacter pylori* peptide deformylase activity. *Biochem Biophys Res Commun* 2013;435:289-94.
41. Baltas N, Karaoglu SA, Tarakci C, Kolayli S. Effect of propolis in gastric disorders: inhibition studies on the growth of *Helicobacter pylori* and production of its urease. *J Enzyme Inhib Med Chem* 2016;31:46-50.
42. Villanueva M, Gonzalez M, Fernandez H, Wilson M, Manquian N, Otth C, et al. In vitro antibacterial activity of Chilean propolis against *Helicobacter pylori*. *Rev Chilena Infectol* 2015;32:530-5.
43. Bergonzelli G, Donnicola D, Porta N, Corthesy-Theulaz I. Essential oils as components of a diet-based approach to management of *Helicobacter* infection. *Antimicrob Agents Chemother* 2003;47:3240-6.
44. Chun SS, Vatter DA, Lin YT, Shetty K. Phenolic antioxidants from clonal oregano (*Origanum vulgare*) with antimicrobial activity against *Helicobacter pylori*. *Process Biochem* 2005;40:809-16.
45. Hawthorne A, Daneshmend T, Hawkey C, Belluzzi A, Everitt S, Holmes G, et al. Treatment of ulcerative colitis with fish oil supplementation: a prospective 12 month randomised controlled trial. *Gut* 1992;33:922-8.
46. Kemen M, Senkal M, Homann HH, Mumme A, Dauphin AK, Baier J, et al. Early postoperative enteral nutrition with arginine-omega-3 fatty acids and ribonucleic acid-supplemented diet versus placebo in cancer patients: An immunologic evaluation of Impact Registered Trademark. *Crit Care Med* 1995;23:652-9.
47. Nordøy A. Is there a rational use for n-3 fatty acids (fish oils) in clinical medicine? *Drugs* 1991;42:331-42.
48. Senkal M, Kemen M, Homann H, Eickhoff U, Baier J, Zumtobel V. Modulation of postoperative immune response by enteral nutrition with a diet enriched with arginine, RNA, and omega-3 fatty acids in patients with upper gastrointestinal cancer. *Eur J Surg* 1995;161:115-22.
49. Thompson L, Cockayne A, Spiller R. Inhibitory effect of polyunsaturated fatty acids on the growth of *Helicobacter pylori*: a possible explanation of the effect of diet on peptic ulceration. *Gut* 1994;35:1557-61.
50. Riber C, Wøjdemann M, Bisgaard T, Ingels H, Rehfeld J, Olsen O. Fish oil reduces gastric acid secretion. *Scand J Gastroenterol* 1999;34:845-8.

S14 Management of multi-drug resistant Helicobacter pylori infection

51. Zurier R. Fatty acids, inflammation and immune responses. *Prostaglandins Leukot Essent Fatty Acids* 1993;48:57-62.
52. Drago L, Mombelli B, Ciardo G, Vecchi ED, Gismondo M. Effects of three different fish Oil7 formulations on *Helicobacter pylori* growth and viability: in vitro study. *J Chemother* 1999;11:207-10.
53. Nualart FJ, Rivas CI, Montecinos VP, Godoy AS, Guaiquil VH, Golde DW, et al. Recycling of vitamin C by a bystander effect. *J Biol Chem* 2003;278:10128-33.
54. Tsao C. An overview of ascorbic acid chemistry and biochemistry. In: Packer L, FuchsJ, eds. *Vitamin C in Health and Disease*. New York: Marcel Dekker 1997;25-58.
55. Deutsch JC. Spontaneous hydrolysis and dehydration of dehydroascorbic acid in aqueous solution. *Anal Biochem* 1998;260:223-9.
56. Sezikli M, Çetinkaya ZA, Sezikli H, Güzelbulut F, Tiftikçi A, Tüzün İnce A, et al. Oxidative stress in *Helicobacter pylori* infection: does supplementation with vitamins C and E increase the eradication rate? *Helicobacter* 2009;14:280-5.
57. Sezikli M, Cetinkaya Z, Güzelbulut F, Yeşil A, Coşgun S, Kurdaş O. Supplementing Vitamins C and E to standard triple therapy for the eradication of *Helicobacter pylori*. *J Clin Pharm and Ther* 2012;37:282-5.
58. Sjunnesson H, Sturegård E, Willén R, Wadström T. High intake of selenium, beta-carotene, and vitamins A, C, and E reduces growth of *Helicobacter pylori* in the guinea Pig. *Comp Med* 2001;51:418-23.
59. Sun YQ, Girgensone I, Leanderson P, Petersson F, Borch K. Effects of antioxidant vitamin supplements on *Helicobacter pylori*-induced gastritis in Mongolian gerbils. *Helicobacter* 2005;10:33-42.
60. Zhang Z, Patchett S, Perrett D, Katelaris P, Domizio P, Farthing M. The relation between gastric vitamin C concentrations, mucosal histology, and CagA seropositivity in the human stomach. *Gut* 1998;43:322-6.
61. Jarosz M, Dzieniszewski J, Dabrowska-Ufniaz E, Wartanowicz M, Ziemiński S, Reed P. Effects of high dose vitamin C treatment on *Helicobacter pylori* infection and total vitamin C concentration in gastric juice. *Eur J Cancer Prev* 1998;7:449-54.
62. Demirci H, Uygun İlikhan S, Öztürk K, Üstündağ Y, Kurt Ö, Bilici M, et al. Influence of vitamin C and E supplementation on the eradication rates of triple and quadruple eradication regimens for *Helicobacter pylori* infection. *Turk J Gastroenterol* 2015;26:456-60.
63. Zojaji H, Talaie R, Mirsattari D, Haghazali M, Molaei M, Mohsenian N, et al. The efficacy of *Helicobacter pylori* eradication regimen with and without vitamin C supplementation. *Dig Liver Dis* 2009;41:644-7.
64. Campanale M, Nucera E, Ojetti V, Cesario V, Di Rienzo T, D'Angelo G, et al. Nickel free-diet enhances the *Helicobacter pylori* eradication rate: a pilot study. *Dig Dis Sci* 2014;59:1851-5.
65. Rostami Nejad M, Rostami K, Yamaoka Y, Mashayekhi R, Molaei M, Dabiri H, et al. Clinical and histological presentation of *Helicobacter pylori* and gluten related gastroenteropathy. *Arch Iran Med* 2011;14:115-18.
66. Minelli M, Schiavino D, Musca F, Bruno M, Falagiani P, Mistrello G, et al. Oral hyposensitization to nickel induces clinical improvement and a decrease in TH1 and TH2 cytokines in patients with systemic nickel allergy syndrome. *Int J Immunopathol Pharmacol* 2010;23:193-201.
67. Fulgione A, Nocerino N, Iannaccone M, Roperto S, Capuano F, Roveri N, et al. Lactoferrin adsorbed onto biomimetic hydroxyapatite nanocrystals controlling-in vivo the *Helicobacter pylori* infection. *PLoS One* 2016;11:0158646.
68. Nazemalhosseini-Mojarad E, Haghghi A, Taghipour N, Keshavarz A, Mohebi SR, Zali MR, et al. Subtype analysis of *Cryptosporidium parvum* and *Cryptosporidium hominis* isolates from humans and cattle in Iran. *Vet Parasitol* 2011;179:250-52.