

**Running head: Irrigation solution temperature and complication of PCNL**

**Effect of Irrigation Solution Temperature on Complication of Percutaneous  
Nephrolithotomy: A Randomized Clinical Trial**

**Seyed Reza Hosseini,<sup>1</sup> Mohammad Ghasem Mohseni,<sup>2</sup> Seyed Mohammad Kazem Aghamir,<sup>3</sup>  
Hamed Rezaei,<sup>4</sup>**

1 Associate Professor, Department of Urology, Tehran University of Medical Sciences ,Tehran, Iran.

Email: srhosseini@tums.ac.ir

2 Professor, Department of Urology, Tehran University of Medical Sciences ,Tehran, Iran. Email:

mgmohseni@tums.ac.ir.

3 Associate Professor, Department of Urology, Tehran University of Medical Sciences, Tehran, Iran.

Email: smkaghamir@yahoo.com

4 Resident of Urology, Tehran University of Medical Sciences ,Tehran, Iran. hamed109@gmail.com

**Key words:** Percutaneous nephrolithotomy, Irrigation fluid, Hypothermia

**Abstract**

**Purpose:** Many factors affecting on hypothermia and shivering during percutaneous nephrolithotomy and in recovery. Hence this study was carried out to determine the effect of irrigation solution temperature on complication of percutaneous nephrolithotomy.

**Materials and Methods:** In this randomized clinical trial, 60 patients under PCNL in Sina University Hospital were enrolled. The patients were randomly assigned in three groups according to simple random manner. The groups included three groups of room temperature fluid (24 degree), warm solution (37 degree), and cold fluid (20 degree) during nephroscopy.

**Results:** Although the initial core temperature was alike across the groups ( $P > 0.05$ ); The hypothermia rate was occurred in all 20 patients in cold fluid group ( $P=0.012$ ). there was significant difference between groups for final temperature and alteration amount ( $P=0.001$ ). The mean VAS scores were significantly lower in warm fluid group compared with the others groups at the recovery, and 8 h postoperatively ( $P = 0.03$ ). Assessment of shivering rates revealed that 3(15%) patients in warm solution group shivered compared with 8 (40%) patients in cold fluid group ( $P=0.018$ ).

**Conclusion:** Warm irrigation solution use during PCNL would result in decrease significantly hypothermia, the mean postoperative pain score and shivering. Hence use of warm irrigation fluid for this matter is recommended.

## **Introduction**

Percutaneous nephrolithotomy (PCNL) is routine treatment for stones larger than two centimeters, staghorn calculi, and resistant to extracorporeal shockwave lithotripsy (ESWL). It is usually done by fluoroscopy or ultrasonography via the calyceal system. Even for the most experienced urologist, major complications can still occur in up to 7% of patients undergoing PNL and minor complications may be encountered in up to 25% of patients. Hemorrhage is the most significant complication of PCNL, with transfusion rates reported to be from less than 1% to 10%. <sup>(1)(17)</sup>

During nephroscopy continuous irrigation of pyelocalyceal system with fluids is required to develop good visual field. The irrigation fluid temperature is studied in different endoscopic procedures showing controversial results. Postoperative hypothermia may result in hazardous complications such as myocardial ischemia, coagulopathy, surgical wound infection, decreased drug metabolism, and

shivering. <sup>(2)</sup>To our knowledge one study about hypothermia in PCNL is available that was evaluated anesthetic complications of hypothermia. <sup>(3)</sup>

The optimal temperature of irrigation solution is not clear and evidence based. Use of warm intra-operative solution may be effective for reduction of postoperative hypothermia risk. On the other hand use of cold fluid may result in better intra-operative homeostasis due to peripheral blood vessels vasoconstriction.

Finding the appropriate temperature for irrigation fluid would result in better surgical outcomes and decreased intra-operative complications such as bleeding and would prepare better visual appearance during nephroscopy, and may result in less hypothermia and related complications. Accordingly, in this study, intra-operative and postoperative complications in PCNL were compared across three groups including those receiving solution with room temperature, warm fluid, and cold solution.

## **Materials and Methods**

### **Study population**

In this randomized clinical trial, 60 patients under PCNL in Sina University Hospital were enrolled. The subjects were in age range of 18 to 60 years. In our department, PCNL is performed in patients with kidney stones more than 2 cm in diameter, stones refractory to extracorporeal shock wave lithotripsy, proximal ureteral stones larger than 1.5 cm in diameter, diverticular stones, and stones producing distal obstruction. The exclusion criteria were medium to high cardiovascular risk, coagulation disorder, renal failure, hepatic failure, diabetes mellitus, and hypothyroidism. The Helsinki Declaration was respected during the study and informed consent form was signed by all patients. Also the study was approved by ethical committee of Tehran University of Medical Sciences.

Patients' enrollment algorithm has been illustrated in Figure 1. Routine Lab. Exam (FBS, CBC, BUN, Creatinine, urine culture) was performed before surgery.

### **Study design**

A randomized multi-arm parallel-group clinical trial with balanced randomization (1:1:1 for three groups) was conducted at the Department of Urology of Tehran University of Medical Sciences in

Tehran, Iran. The study group allocation was by a sequentially running computer-generated block randomization list as blocks of three unique numbers/block, ranging from 1 to 3 unsorted. Sample size was calculated considering a 5 percent expected difference between core temperature in three groups as the primary outcome of interest. we conducted a test with a significance level of 0.05 and power of 0.80 and anticipated that groups of equal size were required. We concluded that at least 20 patients were needed in each group. The groups included three groups of room temperature fluid (24°C), warm solution (37°C), and cold fluid (20°C) during nephroscopy. Demographic data, previous medical history, stone-related data, and operation data were recorded in three groups.

### **Surgical technique**

The patients were not warmed before operation. All patients were transferred to operation room and anesthetized with general method during 20-minute period. The core temperature was assessed and recorded just before initiation of anesthesia. Esophageal temperature probes were planted to measure core temperature. The probe was connected to the monitoring system continuously, during the operations, and was monitored the patients' temperature constantly. However, core temperature recorded its average every 10 minutes. Temperature of the operating room was constantly set at  $23 \pm 1^\circ\text{C}$ , by a central thermostat. Six patients were excluded before operation due to preoperative hypothermia (core temperature less than  $36^\circ\text{C}$ ). The operations were carried out by single practiced surgeon endourologist. The irrigation fluid volume and duration of operation (just prone PCNL time) were also recorded.

After general anesthesia, A 5F urethral catheter was placed cystoscopically and percutaneous access was obtained while the patient was placed in a prone position. Then the access to calyceal system was developed by Shiba needle under fluoroscopy guide and it was dilatated with plastic dilatator up to 30F. Then amplatz sheath was inserted and stones were removed using nephroscope 26F and pneumatic lithoclast. Distilled water was used as irrigation fluid in pressure of 60 mmHg. To ensure patient safety, core temperature was measured during the procedure, if patients suffered severe hypothermia, the surgeon was stopping the irrigation and patient warming using blanket and warmer was being performed.

### ***Outcome assessment***

The core temperature as the primary outcome of interest was again recorded just after operation. The rest of data was Secondary outcomes. Then the patients were transferred to recovery room and underwent routine monitoring for at least one hour. The shivering at recovery room was recorded. Postoperative pain scores were evaluated using a 10-cm self assessed visual analog scale (VAS) with 0 indicating no pain and 10 representing the worst pain experienced by the patient in the recovery room and 8 hours after PCNL.

The reader(fourth author; urology resident) blinded to patient groups. After operation, Routine Lab. exam and plain abdominal radiography were performed. Also, the abdominal CT scan was done as indicated. Stone free was defined as stone diameter less than 4 mm. The complications were categorized to five levels by Modified Clavien system. <sup>(4)</sup>

Data analysis was performed among 60 subjects including three groups. Data analysis was performed by SPSS (version 24.0) software [Statistical Procedures for Social Sciences; Chicago, Illinois, USA]. Fisher exact and Kruskal wallis tests were used and were considered statistically significant at P values less than .05.

### **Results**

Three groups of patients consisting of 20 patients in each were compared. The age, body mass index (BMI), hemoglobin decrease, irrigation volume, stone size, surgical duration, and hospital stay were similar in terms of a number of background variables (Table 1). Although the initial core temperature was alike across the groups ( $P > .05$ ); there was significant difference between groups for final temperature and alteration amount (Tale 2). Eighty percent, 65%, and 70% were male in groups of room temperature fluid, warm solution, and cold fluid, respectively ( $P=.563$ ).

Seventy percent, 80%, and 80% were stone-free in groups of room temperature fluid, warm solution, and cold fluid, respectively ( $P=.700$ ). Assessment of shivering rates revealed that patients in warm solution group shivered less compared with others groups although was not significantly (Table 3) ( $P=.198$ ). The mean VAS scores were significantly lower in warm fluid group compared with the others groups at the recovery, and 8 h postoperatively (Table 3) ( $P = .03$ ). Clavien complications

grading was same across the groups (Table 4). The hypothermia significantly occurred in cold fluid group (Table 5) ( $P=.021$ ).

## **Discussion**

The main finding of the present study is that warm irrigation solution could decrease significantly hypothermia, the mean postoperative pain score and shivering.

Previous studies showed that Cardiovascular, hemorrhagic and infectious complications are significantly more frequent in hypothermic than in normothermic patients. <sup>(2)</sup> Lots of studies have proved that cold stress could influence the immune responses by elevating the levels of inflammatory cytokines, including pro- and anti-inflammatory cytokines. It has been reported that many proinflammatory cytokines, such as TNF- $\alpha$ , IL-1, IL-6, significantly increased under cold stress. For minimally invasive procedure like PCNL, this response is concerned with regional pain. <sup>(5)</sup>

The effects of fluid temperature on core temperature in patients under endoscopic surgeries have been assessed in different studies. The effects on bleeding volume and homeostasis of cold solution are established in some investigations. The effects of experimental lowering of temperature on decreased blood flow are reported by some animal studies. <sup>(6)</sup> Also it has been demonstrated in human studies such as prostatectomy procedures resulting in appropriate hemostasis. <sup>(7)</sup> The bleeding time more than two times after superficial lowering of the temperature is reported in human volunteers. <sup>(8)</sup>

Use of warm irrigation fluid has been also studied in some reports. In the study by Parodi et al. <sup>(9)</sup> use of warm fluid for irrigation during arthroscopy had no effect on reduction hypothermia in shoulder joint but it had significant effect about hip joint. This difference in single study may explain some variations in different studies also. Jin and colleagues <sup>(10)</sup> recommended the use of warm irrigation solution to reduce the hypothermia and shivering and also intra-operative blood loss after endoscopic surgeries; such as our study. Although, in our study hemoglobin level differences were not significant. The isothermal solution led to further fluid overload after operation due to decreased viscosity. <sup>(11)</sup>

There are few study studies in endoscopic urological procedures. Mirza et al reported that hypothermia is common after endoscopic urological procedures which are related to duration of operation, weight, irrigation fluid volume, and type of procedure. <sup>(12)</sup> Rezaei et al showed that using

warm saline irrigation in ureteral endoscopic results in better surgical outcomes including a lower ureteral spasm rate, greater ureteral muscle relaxation and better access to the upper ureteral zone, and a lower rate of complications, such as ureteroscope impaction, ureteral dislodge and stone retropulsion.<sup>(13)</sup> Regarding these confounding factors we matched all these variables across the groups in current study. Also the warm and isothermal irrigation fluids were effective to reduce the hypothermia rate after TURP.<sup>(14)</sup> Use of isothermal fluid was also effective on hypothermia reduction in another study.<sup>(15)</sup> Similar results were also reported by Tekgul and colleagues<sup>(3)</sup> compared the irrigation fluid with room temperature and warm solution in PCNL and reported that lower hypothermia and shivering were seen in warm fluid group. Compare to this study, longer follow up was made in our study during hospitalization to discharge. As well as that study, using warm irrigation fluid resulted in lower hypothermia after procedure, but the complications that could be related to hypothermia, as surgical site infection or coagulopathy was not seen more in patients suffering hypothermia. This may be due to shortness of surgical time or limited number of patients studied in this survey. Actually, longer operation time may exaggerate the impact of irrigation fluid temperature on core body temperature and subsequently such complications. Also in present study, the mean VAS score was significantly lower in the warm fluid group compared with the other groups in the recovery and 8 h postoperatively. In our recent published article was showed that pain score after PCNL has important role to need analgesic drugs.<sup>(16)</sup> Therefore, warm fluid group may be received low dose analgesic drugs compare other groups. Some of limitation of our study were small sample size and limited temperature range of irrigation solutions to compare because of there was no distinct evidence that support to use extreme temperatures in practice . Futher studies is recommended with more patients to attain more reliable results.

## **Conclusions**

Totally, according to our study, it concluded that warm irrigation solution use during PCNL would result in decrease significantly hypothermia, the mean postoperative pain score and shivering. Hence use of warm irrigation fluid for this matter is recommended. However further studies with larger sample size and multi-center sampling are required to attain more definite result with higher reliability and potency for generalization.

## References:

1. Preminger GM, Assimos DG, Lingeman JE, et al. Chapter 1: AUA guideline on management of staghorn calculi: diagnosis and treatment recommendations. *J Urol*. 2005;173:1991–2000.
2. Torossian A, Brauer A, Hocker J, Bein B, Wulf H, Horn EP. Preventing inadvertent perioperative hypothermia. *Deutsches Arzteblatt international*. 2015; 112:166-72.
3. Tekgul ZT, Pektas S, Yildirim U, et al. A prospective randomized double-blind study on the effects of the temperature of irrigation solutions on thermoregulation and postoperative complications in percutaneous nephrolithotomy. *J Anesth*. 2015; 29:165-9.
4. Tefekli A, Ali Karadag M, Tepeler K, et al. Classification of percutaneous nephrolithotomy complications using the modified claviengrading system: looking for a standard. *Eur Urol*. 2008; 53:184-90.
5. Guo JR, Li SZ, Fang HG. Different duration of cold stress enhances pro-inflammatory cytokines profile and alterations of Th1 and Th2 type cytokines secretion in serum of wistar rats. *J Anim Vet Adv*. 2012; 11:1538–1545
6. Venjakob AJ, Vogt S, Stockl K, Tischer T, Jost PJ, Thein E. Local cooling reduces regional bone blood flow. *Journal of orthopaedic research: official publication of the Orthopaedic Research Society*. 2013; 31:1820-7.
7. Zorn KC, Bhojani N, Gautam G, Shikanov S, Gofrit ON, Jayram G. Application of ice cold irrigation during vascular pedicle control of robot-assisted radical prostatectomy: EnSeal instrument cooling to reduce collateral thermal tissue damage. *J Endourol*. 2010 24:1991-6.
8. Romlin B, Petruson K, Nilsson K. Moderate superficial hypothermia prolongs bleeding time in humans. *Acta Anaesthesiol Scand*. 2007;51:198-201.
9. Parodi D, Valderrama J, Tobar C, Besomi J, Lopez J, Lara J. Effect of warmed irrigation solution on core body temperature during hip arthroscopy for femoroacetabular impingement. *Arthroscopy: the journal of arthroscopic & related surgery: official publication*



- of the Arthroscopy Association of North America and the International Arthroscopy Association. 2014; 30: 36-41.
10. Jin Y, Tian J, Sun M, Yang KA. systematic review of randomised controlled trials of the effects of warmed irrigation fluid on core body temperature during endoscopic surgeries. *J Clin Nurs*. 2011; 20:305-16.
  11. Kim YS, Lee JY, Yang SC, Song JH, Koh HS, Park WK. Comparative study of the influence of room-temperature and warmed fluid irrigation on body temperature in arthroscopic shoulder surgery. *Arthroscopy: the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association*. 2009; 25:24-9.
  12. Mirza S, Panesar S, AuYong KJ, French J, Jones D, Akmal S. The effects of irrigation fluid on core temperature in endoscopic urological surgery. *J Perioper Pract* 2007; 17:49-51.
  13. Mohammadzadeh Rezaei MA, Akhavan Rezayat A, Tavakoli M, Jarahi L. Evaluation the result of warm normal saline irrigation in ureteral endoscopic surgeries. *Urol J*. 2018; 15:83-86.
  14. Okeke LI. Effect of warm intravenous and irrigating fluids on body temperature during transurethral resection of the prostate gland. *BMC urol*. 2007; 7:15.
  15. Pit MJ, Tegelaar RJ, Venema PL. Isothermic irrigation during transurethral resection of the prostate: effects on peri-operative hypothermia, blood loss, resection time and patient satisfaction. *Br J Urol*. 1996; 78:99-103.
  16. Hosseini SR, Imani F, Shayanpour G, Khajavi MR. The effect of nephrostomy tract infiltration of ketamine on postoperative pain and peak expiratory flow rate in patients undergoing tubeless percutaneous nephrolithotomy: a prospective randomized clinical trial. *Urolithiasis*. 2017 ;45:591-595.
  17. Maghsoudi R, Etemadian M, Kashi AH, Ranjbaran A. The Association of Stone Opacity in Plain Radiography with Percutaneous Nephrolithotomy Outcomes and Complications. *Urol J*. 2016 8;13:2899-2902.

**Corresponding Author:** Seyed Reza Hosseini; **Email:** rhosseinim@yahoo.com

**Mailing Address:** Department of Urology, Sina Hospital, Imam Khomeini Ave., Tehran, Iran

**Table 1- Background data across the groups**

<b>Variable</b>	<b>Room temperature fluid</b>	<b>Body temperature fluid</b>	<b>Cold fluid</b>	<b>P Value</b>
<b>Age</b>	45.9 ± 11.9	44.9 ± 12.2	39.9 ± 16.6	0.345
<b>BMI (kg/m<sup>2</sup>)</b>	26.5 ± 3.4	25.6 ± 2.1	24.8 ± 3.6	0.241
<b>hemoglobin decrease (g/dl)</b>	1.6 ± 1.1	1.5 ± 0.9	2.1 ± 1.0	0.132
<b>irrigation volume (liter)</b>	12.1 ± 2.8	12.6 ± 3.3	12.8 ± 3.2	0.768
<b>stone size (cm)</b>	3.1 ± 0.9	3.2 ± 1.1	3.6 ± 1.3	0.331
<b>Operation duration (min)</b>	85.0 ± 31.2	85.7 ± 38.7	86.5 ± 30.1	0.988
<b>Hospital stay (day)</b>	5.2 ± 2.5	4.8 ± 1.4	5.1 ± 2.3	0.778
<b>Creatinine increase(mg/dl)</b>	0.5±0.2	0.4±0.1	0.6±0.4	0.11

**Table 2- Core temperature across the groups**

<b>Variable</b>	<b>Room temperature fluid</b>	<b>Body temperature fluid</b>	<b>Cold fluid</b>	<b>P Value</b>
<b>Initial temperature</b>	36.8 ± 0.4	36.7 ± 0.4	36.6 ± .4	0.259
<b>Final temperature</b>	35.7 ± 0.9	36.1 ± 0.6	35.0 ± 1.1	0.001
<b>Temperature alteration</b>	1.1 ± 0.8	0.6 ± 0.4	1.6 ± 0.9	0.001

**Table 3- Complications across the groups**


<b>Variable</b>	<b>Room temperature fluid</b>	<b>Body temperature fluid</b>	<b>Cold fluid</b>	<b>p-value</b>	<b>test</b>
<b>Shivering*</b>	5 (25%)	3 (15%)	8 (40%)	0.198	
<b>Fever</b>	1 (5%)	3 (15%)	3 (15%)	0.68	Fisher exact
<b>DVT</b>	---	---	1 (5%)	1.000	Fisher exact
<b>Angioembolization</b>	1 (5%)	---	---	1.000	Fisher exact
<b>Transfusion</b>	1 (5%)	1 (5%)	1 (5%)	1.000	
...					

**Table 4- Clavein complications grading across the groups**

<b>Grade</b>	<b>Room temperature fluid</b>	<b>Body temperature fluid</b>	<b>Cold fluid</b>	<b>p-value of kruskal wallis</b>
<b>1</b>	2 (10%)	1 (5%)	1 (5%)	0.910
<b>2</b>	2 (10%)	3 (15%)	4 (20%)	
<b>3</b>	1 (5%)	---	---	
<b>Negative</b>	15 (75%)	16 (80%)	15 (75%)	

**Table 5- Hypothermia across the groups**

<b>Grade</b>	<b>Mild (34-36 degree)</b>	<b>Moderate (32-33.9 degree)</b>	<b>Negative</b>	<b>True p-value of kruskal wallis</b>
<b>Room temperature fluid</b>	11 (55%)	1 (5%)	8 (40%)	0.021
<b>Body temperature fluid</b>	6 (30%)	---	14 (70%)	
<b>Cold fluid</b>	13 (65%)	3 (15%)	4 (20%)	



**Figure 1.** Patient flow chart of the clinical study