

Nutritional and Phytochemical Profiling of African Star Apple (*Chrysophyllum albidum*) and Lagos Spinach (*Celosia argentea*): Unlocking Their Potential for Human Health

Hassan Abdulsalam Adewuyi^{1*}, Lukman Adeola Usman², Fatima Mahmoud Muhammad¹, Oyewola Abdulkabir³, Mba Michael Okechukwu⁴, Moses Orimoloye Akinjiyan⁵, Adebimpe Hameedah Oluwatoyin⁶, Khairat Shamsudeen⁵

¹. Department of Biochemistry, Federal University of Technology, PMB 65 Minna, Niger State Nigeria

². Department of Biological Sciences Ahman Pategi University, Pategi Kwara State Nigeria

³. Department of Fisheries Lagos State University, Lagos Nigeria

⁴. Department of Biochemistry, Faculty of Life Sciences, University of Benin, Benin City Nigeria

⁵. Department of Medical Biochemistry, School of Basic Medical Sciences, Federal University of Technology, Akure

⁶. Department of Chemistry and Biochemistry, College of Health and Natural Sciences-Keplinger Hall, The University of Tulsa, Oklahoma, USA, 74104

⁷. Department of Microbiology, Saadu Zungur University, Bauchi State Nigeria



Cite this article as: Adewuyi, H. A., Usman, L. A., Muhammad, F. M., Abdulkabir, O., Okechukwu, M. M., Akinjiyan, M. O., ... Shamsudeen, K. (2025). Nutritional and Phytochemical Profiling of African Star Apple (*Chrysophyllum albidum*) and Lagos Spinach (*Celosia argentea*): Unlocking Their Potential for Human Health. *Archives of Advances in Biosciences*, 16(1), 1–6. <https://doi.org/10.22037/aab.v16i1.50357>

<https://journals.sbmu.ac.ir/aab/article/view/50357>



Article info:

Received: 20 Sep 2025

Accepted: 03 Nov 2025

Published: 17 Nov 2025

* Corresponding author:

Hassan Abdulsalam Adewuyi, MSc.

Address: Department of Biochemistry, Federal University of Technology, PMB 65 Minna, Niger State Nigeria

E-mail: :

hasselbatch@gmail.com

Abstract

Introduction: Indigenous African plants are increasingly studied as sustainable sources of nutrients and bioactive compounds. *Chrysophyllum albidum* (African star apple) and *Celosia argentea* (Lagos spinach) are widely consumed as food and medicine, yet their biomedical potential remains insufficiently explored. This study was designed to comprehensively evaluate their nutritional composition, phytochemical profile, antioxidant capacity, and antimicrobial properties.

Materials and Methods: Ripe fruits of *C. albidum* and leaves of *C. argentea* were collected and authenticated. Proximate and micronutrient analyses were conducted using AOAC methods. Phytochemicals were quantified via HPLC and GC-MS. Antioxidant capacity was assessed by DPPH and FRAP assays. Antimicrobial activity against bacteria and fungi was determined by MIC and MBC.

Results: *C. albidum* had significantly higher levels of carbohydrate ($56.3 \pm 1.2\%$) and vitamin C (41.8 ± 2.1 mg/100 g), while *C. argentea* was richer in protein ($23.6 \pm 0.9\%$), calcium (280.7 ± 6.3 mg/100 g), and iron (18.2 ± 0.7 mg/100 g). Phytochemical profiling revealed substantial flavonoids, phenolics, saponins, terpenoids, alkaloids, and carotenoids, with *C. argentea* showing significantly higher flavonoids and carotenoids contents. Antioxidant assays indicated strong radical scavenging activity (DPPH IC₅₀: *C. albidum* 42.6 µg/mL; *C. argentea* 38.2 µg/mL). Both extracts inhibited microbial growth, with MIC values as low as 125 µg/mL against *Staphylococcus aureus*.

Conclusion: Both plants exhibit nutritional and phytochemical richness, potent antioxidant capacity, and antimicrobial potential. Their integration into diets and nutraceutical development could aid in reducing oxidative stress and microbial infections.

Keywords: *Chrysophyllum albidum*; *Celosia argentea*; phytochemicals; antioxidant; antimicrobial; nutrition.

1. Introduction

Functional foods derived from plants are gaining attention as safe and sustainable strategies to prevent nutrition-related disorders and chronic diseases [1]. In sub-Saharan Africa, indigenous plants serve as both dietary staples and medicinal remedies, offering dual nutritional and therapeutic benefits [2]. Among such plants, *Chrysophyllum albidum* (African star apple) and *Celosia argentea* (Lagos spinach) are particularly notable for their widespread and traditional use.

C. albidum, a member of the Sapotaceae family, produces fruits consumed during the dry season. The fruit pulp is known for its sweet-sour taste and its richness in sugar, vitamins, and minerals [3]. Traditionally, it has been used to manage malaria, wounds, and infections. *C. argentea*, a leafy vegetable of the Amaranthaceae family, is cultivated across West Africa. Its leaves are consumed in soups or sauces. Traditional healers also use it for anemia, gastrointestinal disorders, and reproductive health [4,5].

The nutritional profiling of these plants is important for establishing their contributions to human health. Proximate and micronutrient analyses identify their energy and mineral values, while phytochemical profiling reveals secondary metabolites, such as flavonoids and phenolics, that provide antioxidant and antimicrobial activities [6–8]. Studies have linked plant-derived antioxidants to the reduction of oxidative stress-related conditions such as cardiovascular disease, cancer, and diabetes [9]. Similarly, phytochemicals with antimicrobial activity can provide alternatives to synthetic drugs, especially in the context of antimicrobial resistance [10].

Although some studies have examined these plants individually, a comparative evaluation of their nutritional, phytochemical, antioxidant, and antimicrobial profiles is lacking. This study bridges this gap by comprehensively analyzing *C. albidum* fruits and *C. argentea* leaves. The findings will contribute to functional food research, nutraceutical development, and public health nutrition.

2. Materials and Methods

Sample Collection and Authentication

Ripe *C. albidum* fruits and fresh *C. argentea* leaves were purchased from a farm market in Southwestern Nigeria. The plant samples were authenticated and

voucher specimens were deposited at the Department of Botany, Saadu Zungur University, Bauchi.

Proximate and Micronutrient Analysis

Moisture, ash, protein, fat, fiber, and carbohydrate contents were determined using AOAC standard methods [11]. Micronutrient content was analyzed as follows: iron, calcium, and potassium by atomic absorption spectrometry; vitamin C by titrimetry; and vitamin A by spectrophotometry.

Phytochemical Analysis

Qualitative screening of phytochemicals (alkaloids, flavonoids, tannins, phenolics, saponins, terpenoids, and carotenoids) was performed using standard methods. Quantitative analysis of flavonoids and phenolics was performed by HPLC, while carotenoids were estimated by UV-Vis spectrophotometry. GC-MS was used for metabolite fingerprinting.

Antioxidant Assays

The antioxidant potential of the extracts was assessed using the DPPH radical scavenging assay and the FRAP (ferric reducing antioxidant power) assay. For the DPPH, a 1 mL aliquot of extract at different concentrations (25–200 µg/mL) was mixed with 1 mL of 0.1 mM DPPH solution in methanol. The mixture was incubated in the dark at room temperature for 30 minutes, and absorbance was measured at 517 nm. The percentage scavenging activity was calculated using the control absorbance, and IC₅₀ values were determined. The FRAP Assay, a 100 µL extract sample was reacted with 900 µL of FRAP reagent (composed of 300 mM acetate buffer, 10 mM TPTZ, and 20 mM FeCl₃·6H₂O). The mixture was incubated for 10 minutes at 37°C, and absorbance was recorded at 593 nm. Results were expressed as µmol ascorbic acid equivalent per gram (µmol AAE/g).

Antimicrobial Assays

Antimicrobial activity was evaluated using the broth microdilution technique following the method described by [12]. The tested microorganisms included *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Candida albicans*. Microbial inocula were standardized to 0.5 McFarland turbidity (~1.5 × 10⁸ CFU/mL). Serial dilutions of the plant extracts (1000–62.5 µg/mL) were prepared in sterile 96-well microplates. Following inoculation, the plates were incubated at 37°C for 24 hours for bacteria and at 28°C for 48 hours for fungi. The minimum

inhibitory concentration (MIC) was determined as the lowest concentration showing no visible growth, while the minimum bactericidal concentration (MBC) or minimum fungicidal concentration (MFC) was established by subculturing aliquots from non-turbid wells onto nutrient agar plates to confirm microbial death.

Statistical Analysis

All analyses were performed in triplicate. Results are presented as mean \pm standard deviation. Statistical comparisons were conducted using One-way

ANOVA, followed by Tukey's test; $p < 0.05$ was considered statistically significant.

3. Results

Proximate composition of *C. albidum* and *C. argentea*

C. argentea was significantly higher in protein, fat, fiber, and ash ($p < 0.05$), confirming its value as a nutrient-dense vegetable for protein and mineral supplementation. *C. albidum* contained significantly higher carbohydrates ($p < 0.05$), making it a high-energy fruit. These differences suggest complementary nutritional roles.

Table 1. Proximate composition of *C. albidum* and *C. argentea*

Parameter	<i>C. albidum</i> (%)	<i>C. argentea</i> (%)
Moisture	8.4 \pm 0.5 ^b	9.6 \pm 0.4 ^a
Ash	3.1 \pm 0.2 ^b	3.8 \pm 0.1 ^a
Protein	7.9 \pm 0.3 ^b	23.6 \pm 0.9 ^a
Fat	2.8 \pm 0.1 ^b	4.2 \pm 0.2 ^a
Fiber	4.6 \pm 0.2 ^b	5.4 \pm 0.3 ^a
Carbohydrates	56.3 \pm 1.2 ^a	49.2 \pm 1.0 ^b

Values are expressed as mean \pm SD (n = 5). Values with different superscript letters (a, b, c) indicate statistically significant differences at $p < 0.05$.

Micronutrient composition of *C. albidum* and *C. argentea*

C. albidum had significantly higher vitamin C and potassium ($p < 0.05$), enhancing immune defense and

electrolyte regulation. *C. argentea* was significantly richer in vitamin A, calcium, and iron ($p < 0.05$), indicating strong potential for preventing anemia and bone disorders.

Table 2. Micronutrient composition of *C. albidum* and *C. argentea*

Nutrient	<i>C. albidum</i>	<i>C. argentea</i>
Vitamin C (mg/100g)	41.8 \pm 2.1 ^a	36.2 \pm 1.8 ^b
Vitamin A (μ g/100g)	225.4 \pm 7.3 ^b	312.1 \pm 6.5 ^a
Iron (mg/100g)	12.4 \pm 0.5 ^b	18.2 \pm 0.7 ^a
Calcium (mg/100g)	198.5 \pm 5.2 ^b	280.7 \pm 6.3 ^a

Potassium (mg/100g)	312.4 ± 8.6 ^a	290.3 ± 7.4 ^b
---------------------	--------------------------	--------------------------

Values are expressed as mean ± SD (n = 5). Values with different superscript letters (a, b, c) indicate statistically significant differences at p < 0.05.

Phytochemical composition of *C. albidum* and *C. argentea*

C. argentea exhibited significantly higher levels of flavonoids, saponins, terpenoids, alkaloids, and

carotenoids (p < 0.05), conferring superior therapeutic potential. Conversely, *C. albidum* showed significantly higher phenolics (p < 0.05), consistent with its strong antioxidant properties.

Table 3. Phytochemical composition of *C. albidum* and *C. argentea*

Compound	<i>C. albidum</i>	<i>C. argentea</i>
Flavonoids	45.2 ± 1.7 ^b	62.5 ± 2.3 ^a
Phenolics	89.6 ± 3.2 ^a	78.4 ± 2.9 ^b
Saponins	23.1 ± 0.9 ^b	28.9 ± 1.2 ^a
Terpenoids	14.3 ± 0.6 ^b	19.2 ± 0.8 ^a
Alkaloids	11.5 ± 0.5 ^b	15.7 ± 0.6 ^a
Carotenoids	6.8 ± 0.3 ^b	9.5 ± 0.4 ^a

Values are expressed as mean ± SD (n = 5). Values with different superscript letters (a, b, c) indicate statistically significant differences at p < 0.05.

Antioxidant activity of *C. albidum* and *C. argentea*

Both species exhibited strong antioxidant activities. *C. argentea* demonstrated significantly stronger

scavenging and reducing power (p < 0.05), suggesting superior potential for oxidative stress reduction.

Table 4. Antioxidant activity of *C. albidum* and *C. argentea*

Assay	<i>C. albidum</i>	<i>C. argentea</i>
DPPH IC ₅₀ (µg/mL)	42.6 ± 1.8 ^b	38.2 ± 1.6 ^a
FRAP (µmol AAE/g)	812.4 ± 25.7 ^b	924.5 ± 28.3 ^a

Values are expressed as mean ± SD (n = 5). Values with different superscript letters (a, b, c) indicate statistically significant differences at p < 0.05.

Antimicrobial activity of *C. albidum* and *C. argentea*

Both plants exhibited a notable antimicrobial activity. MIC values were lowest against *S.*

aureus and *C. albicans* (125–250 µg/mL), confirming their traditional use against infections.

Table 5. Antimicrobial activity of *C. albidum* and *C. argentea*

Microorganism	<i>C. albidum</i> (µg/mL)	<i>C. argentea</i> (µg/mL)
<i>Staphylococcus aureus</i>	125	125

<i>Escherichia coli</i>	250	250
<i>Pseudomonas aeruginosa</i>	500	500
<i>Candida albicans</i>	250	250

4. Discussion

This study provides a comparative analysis of the nutritional and phytochemical profiles of *C. albidum* and *C. argentea*. The proximate data show that *C. argentea* is rich in protein, consistent with previous studies identifying African green leafy vegetables as affordable protein supplements [13]. In contrast, *C. albidum*'s carbohydrate content supports its role as an energy fruit.

Micronutrient data reveals that *C. albidum* is a fruit rich in vitamin C and potassium. This finding aligns with its traditional use in enhancing immunity and fluid balance [14]. Conversely, *C. argentea*'s iron and calcium content highlight its potential in preventing anemia and improving bone health [15].

Phytochemical results demonstrate that *C. argentea* contains higher levels of flavonoids, carotenoids, and alkaloids, compounds known for modulating inflammation and protecting against oxidative damage [16]. On the other hand, *C. albidum*'s higher phenolic content confirms its antioxidant potential, in agreement with findings from other tropical fruits [17].

Antioxidant assays confirm potent radical scavenging and reducing capacities in both species, particularly in *C. argentea* which can explain its cardioprotective and hepatoprotective properties as documented in traditional medicine [18,19].

Finally, both species exhibited antimicrobial activity, especially against *S. aureus* and *C. albicans*. These results support previous studies on African botanicals as sources of antimicrobial agents, particularly for managing resistant strains [20,21].

Collectively, these findings demonstrate the functional food potential of *C. albidum* and *C. argentea*. Their integration into dietary practices and nutraceutical formulations could help address nutritional deficiencies, oxidative stress, and infectious diseases in African populations and beyond.

5. Conclusion

Chrysophyllum albidum and *Celosia argentea* are nutrient-rich and phytochemically diverse, exhibiting strong antioxidant and antimicrobial properties. Their

consumption, along with further research into their bioactive compounds, is recommended for the development of functional foods and nutraceuticals..

Ethical Considerations

Compliance with ethical guidelines

The study did not involve human or animal subjects. Ethical approval was therefore not required.

Funding

This research received no external funding.

Author's contributions

Conceptualization: Hassan Abdulsalam Adewuyi; Methodology: Khairat Shamsudeen; Data Analysis: Lukman Adeola Usman; Writing – Original Draft: Oyewola Abdulkabir; Writing – Review and Editing: All authors; Supervision: Fatima Mahmoud Muhammad; Data Curation and Analysis: Mba Michael Okechukwu; Adebimpe Hameedah Oluwatoyin: Data Curation and Analysis

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

The authors thank the Department of Biochemistry, Saadu Zungur University Bauchi State Nigeria, for providing laboratory support.

References

- [1] Yusuf AA, Lawal B, Alozieuwa UB, Onikanni AS, Lukman HY, Fadaka AO, Olawale F, Osuji O, Sani S, Owolabi MS, Adewuyi AH. Attenuating effects of *Azanza garckeana* fractions on glycemo-impaired-associated dyslipidemia, hepatopathy, and nephropathy. *American journal of translational research*. 2023 Oct 15;15(10):5997. [\[PMCID\]](#)
- [2] Adewuyi HA, Kabiru AY, Muhammad HL, Lukman HY, Owolabi MS, Jonathan I, El-Gazzar AM, Mahmoud MH, Batiha GE, Lawal B. Pre-clinical protective potentials of *Carica papaya* constituents in experimentally induced anemia.

- American Journal of Translational Research. 2024 Jul 15;16(7):3259. [\[PMID\]](#)
- [3] Fagbohun BO, Adewuyi HA, Musa M, Unuata ON, Ogar OE, Achi AE, Mohammed UA, Olukayode JT. Evaluation of Albizia lebbek and Curcuma longa Extracts on Gastrointestinal Motility, Safety, and Hematological Parameters: A Sub-Chronic Toxicity and Pharmacological Study. Asia Pacific Journal of Medical Toxicology. 2025 Mar 1;14(1). [\[LINK\]](#)
- [4] Hastata L. Protective Role of Leptadenia Hastata on the Haematological and Biochemical Alterations in Adrenaline-Induced Hypertensive Rats. [\[LINK\]](#)
- [5] OGAR OE, MESSIAH AI, ADEWUYI HA, ADIO WS, OLUWATOYIN AH, OBUNADIKE CV, MOHAMMED UA, SHEKARI A, AUGUSTINE B, SEBASTINE KL, OLUKAYODE JT. Anti-hyperglycemic, Anti-hyperlipidemic, Hematological, Hepatoprotective, And Antioxidant Effects of Vernonia amygdalina on Alloxan-induced Diabetic Wistar Rats. [\[LINK\]](#)
- [6] Adewuyi HA, Shekari A, Adio SW, Oluwatoyin AH, Fagbohun RO, Owwoye EA, Korie GC, Nasiru MO, Olusegun TG, Ishola AB, Gidado MO. Ameliorative Effects of Bridelia ferruginea Extracts on Cadmium Chloride-Induced Reproductive Hormone Imbalance, Oxidative Stress, Hepatorenal Damage, Hematological Disorders, and Acute Toxicity in Wistar Rats. bioRxiv. 2025 Jan 15:2025-01. [\[LINK\]](#)
- [7] Fagbohun BO, Unuata ON, Enogela AB, Adewuyi HA, Jonathan I, Gyang DD, Adio WS, Oluwatoyin AH, Mohamed UA, Ogar OE. Phytochemical Characterization, Toxicological Evaluation, and Biological Activities of Phenolics Fractions from Chromolaena Odorata: A Subacute Toxicity Study. [\[LINK\]](#)
- [8] Adewuyi HA, Ugwu CS, Adio SW, Nasiru MO, Kolawole AV, Adeniji AE, Tairu SK, Jumah TA, Adekoya AM, Olaniran TT, Adeniyi EO. Chemotherapeutic potential of methanol leaf extract of Telfairia occidentalis on oxidative stress, hepatic, hematological, and biochemical alterations in DMBA-induced breast cancer in Wistar rats. Archives of Advances in Biosciences. 2025;16(1):1-7. [\[LINK\]](#)
- [9] Adewuyi HA, Atapia M, Ochonung E, Waheed SA, Oluwatoyin AH, Obunadike CV, Edema AA, Hussein S, Mohammed UA, Sebastine KL, Jaiyeola TO. Phytochemical and nutritional composition of Vernonia amygdalina: antioxidant, antidiabetic efficacy, and hematological benefits. Pharmacology and Clinical Pharmacy Research. 2025;10(2). [\[LINK\]](#)
- [10] Bushirat FO, Adewuyi HA, Adio SW, Musa MN, Olusegun TG, Ishola AB, Idoko A, Kolawole AV, Oluwatoyin AA, Agwasim SN. Phytochemical analysis, antioxidant, and antibacterial properties of partition fractions of Adansonia digitata and Annona muricata extracts using chloroform, ethyl acetate, ethanol, and aqueous solvent systems. Journal of Biomedical and Clinical Research. 2025 Jul 24;18:199-213. [\[LINK\]](#)
- [11] Fagbohun BO, Ogar OE, Messiah AI, Adewuyi HA, Adio WS, Oluwatoyin AH, Obunadike CV, Edema AA, Hussein S, Mohammed UA, Jaiyeola TO. Phytotherapeutic approach to hypertension: a randomized control trial of Euphorbia hirta and Leptadenia hastata extracts in adrenaline-induced hypertensive Wistar rats. Pharm Biomed Res. 2025;11(2):3-10. [\[LINK\]](#)
- [12] AOAC. Official Methods of Analysis. 21st ed. Gaithersburg, MD: AOAC International; 2019.
- [13] Odhav B, Beekrum S, Akula US, Baijnath H. Preliminary assessment of nutritional value of traditional leafy vegetables in KwaZulu-Natal, South Africa. Journal of Food Composition and Analysis. 2007 Aug 1;20(5):430-5. [\[LINK\]](#)
- [14] Akinmoladun FO, Komolafe TR, Komolafe EO. Phytochemical screening and proximate analysis of Chrysophyllum albidum. J Appl Sci Environ Manag. 2017;21(5):1037-41. [\[Doi:10.4314/jasem.v21i5.31\]](#)
- [15] Ogunlade I, Olaofe O, Fadare DA. Nutritional potential of the seed and pulp of African star apple (Chrysophyllum albidum). J Food Technol. 2005;3(4):345-8. [\[LINK\]](#)
- [16] Ndlovu J, Afolayan AJ. Nutritional analysis of the South African wild vegetable Corchorus olitorius L. Asian J. Plant Sci. 2008 Nov 5;7(6):615-8. [\[LINK\]](#)
- [17] Falodun A, Nworgu ZAM, Ikponmwosa E, Adesanya SA. Phytochemical screening and in vitro antioxidant activities of Celosia argentea. Sci World J. 2010;5(2):5-9.

- [18] Adetuyi AO, Osagie AU. Nutrient, antinutrient and phytochemical composition of *Celosia argentea* leaves. *Niger J Basic Appl Sci.* 2011;19(1):81-4. [\[LINK\]](#)
- [19] Nguyen HC, Chen CC, Lin KH, Chao PY, Lin HH, Huang MY. Bioactive compounds, antioxidants, and health benefits of sweet potato leaves. *Molecules.* 2021 Mar 24;26(7):1820. [\[LINK\]](#)
- [20] Akinmoladun FO, Komolafe TR, Farombi EO. Antioxidant and antimicrobial activities of *Chrysophyllum albidum* fruit extracts. *J Med Food.* 2019;22(2):212-9. [\[Doi:10.1089/jmf.2018.0086\]](#)
- [21] Oboh G, Ademiluyi AO, Ademosun AO. Phenolic extract from *Celosia argentea* leaves inhibits key enzymes linked to type 2 diabetes and hypertension. *J Evid Based Complementary Altern Med.* 2013;18(1):47-54. [\[Doi:10.1177/2156587212465077\]](#)