

REVIEW ARTICLE

Rapid Arterial Occlusion Evaluation (RACE) Tool in Detecting Large Cerebral Vessel Occlusions; a Systematic Review and Meta-Analysis

Iman Chehregani Rad¹, Amir Azimi^{2*}

1. Physiology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Rajaie Cardiovascular Medical and Research Center, Iran university of medical sciences, Tehran, Iran

Received: November 2023; Accepted: December 2023; Published online: 14 November 2023

Abstract: **Introduction:** Large vessel occlusion (LVO) strokes are linked to higher mortality rates and a greater risk of long-term disability. This study aimed to evaluate the diagnostic performance of the Rapid Arterial Occlusion Evaluation (RACE) tool in detecting LVO through a systematic review and meta-analysis. **Methods:** A comprehensive search was conducted across online databases including PubMed, Embase, Scopus, and Web of Science, up to June 25th, 2023. Additionally, a manual search on Google and Google Scholar was performed to identify studies that assessed the diagnostic accuracy of the RACE scale in detecting LVO among patients with stroke symptoms. **Results:** Data extracted from 43 studies were analyzed. The optimal cut-off points were determined to be 3 and 4, with a sensitivity of 0.86 (95% confidence interval (CI): 0.78, 0.91) and specificity of 0.57 (95% CI: 0.49, 0.67) for cut-off ≥ 3 , and a sensitivity of 0.78 (95% CI: 0.70, 0.84) and specificity of 0.68 (95% CI: 0.59, 0.75) for cut-off ≥ 4 . Subgroup meta-regression analysis revealed significant variations in sensitivity and specificity. RACE scale's sensitivity was significantly higher in LVO detection in suspected stroke cases, in pre-hospital settings, prospective design studies, and when considering both anterior and posterior occlusions for LVO definition. RACE scale's specificity was significantly higher when evaluating confirmed stroke cases, in-hospital settings, and considering only anterior occlusions for LVO definition and retrospective design studies. Notably, RACE exhibited higher sensitivity and specificity when utilized by neurologists and physicians compared to other emergency staff. Despite these variations, our study found comparable diagnostic accuracy across different conditions. **Conclusion:** A high level of evidence indicates that the RACE scale lacks promising diagnostic value for detection of LVOs. A sensitivity range of 0.69 to 0.86 is insufficient for a screening tool intended to aid in the diagnosis of strokes, considering the substantial morbidity and mortality associated with this condition.

Keywords: Arterial occlusive diseases; Clinical decision rules; Diagnosis; Systematic review; Meta-analysis

Cite this article as: Chehregani Rad I, Azimi A. Rapid Arterial Occlusion Evaluation (RACE) Tool in Detecting Large Cerebral Vessel Occlusions; a Systematic Review and Meta-Analysis. Arch Acad Emerg Med. 2024; 12(1): e10. <https://doi.org/10.22037/aaem.v12i1.2152>.

1. Introduction

Large vessel occlusion (LVO) is a critical condition characterized by the blockage of major cerebral arteries, accounting for 24-46% of acute ischemic strokes (AIS). Compared to non-LVO ischemic strokes, LVO strokes are linked to higher mortality rates and a greater risk of long-term disability (1, 2). Notably, a meta-analysis revealed that LVO patients had a 64% incidence of unfavorable outcomes and a mortality rate of 26.2%, in contrast to the 24% poor outcomes and 1.3% mortality rate observed in non-LVO cases (2).

Endovascular treatment (EVT) has emerged as the gold stan-

dard in management of LVO (3). Guidelines strongly recommend initiating EVT within 6 hours from the patient's last known well time (4). Studies suggest that even within a window of 6 to 16 hours since the last known well time, LVO patients can benefit from EVT (5). Some studies have even extended this treatment window to 6 to 24 hours (3).

However, not all healthcare facilities have the capability to perform EVT. Thus, it becomes imperative to rapidly identify LVO cases and bypass primary emergency centers, directing patients to Comprehensive Stroke Centers (CSCs) to ensure they don't miss the optimal treatment window (6).

Diagnosing LVO typically relies on advanced imaging techniques such as magnetic resonance imaging (MRI) and computed tomography (CT) scans (1). Nonetheless, specialized scales have been designed for rapid and accurate LVO diagnosis, including, FAST-ED (Face-Arm-Speech-Time-Eye-Disturbance), VAN (Vision-Aphasia-Neglect), PASS (Prehos-

*Corresponding Author: Amir Azimi; Rajaie Cardiovascular Medical and Research Center, Hashemi rafsanjani, Tehran, Iran. Postal Code: 9619763142. Tel: +989123957125, Email: azimi7619@gmail.com. ORCID: <https://orcid.org/0000-0002-8317-7009>.

pital Acute Stroke Severity), LEGS (Lever Arm Test-Gaze-Eyes-Visual Field-Speech), CPSSS (Cincinnati Prehospital Stroke Severity Scale), LAMS (Los Angeles Motor Scale), RACE (Rapid Arterial Occlusion Evaluation), and NIHSS (National Institutes of Health Stroke Scale) (7-14). Among these, the RACE scale stands out for its simplicity and rapid assessment. The RACE scale comprises five essential components: assessment of facial palsy, arm motor function, leg motor function, head and gaze deviation, as well as aphasia and agnosia, with scores ranging from 0 to 9 (7).

While a previous study has systematically evaluated the RACE scale's performance (15), its clinical utility, applicability in diverse settings, with varying assessors, and under various study designs remain underexplored. Furthermore, the previous systematic review is outdated and it is essential to incorporate the findings of the many newly published studies. Therefore, the objective of this systematic review and meta-analysis was to assess the diagnostic accuracy of the Rapid Arterial Occlusion Evaluation (RACE) scale using different cut-off values and to ascertain its effectiveness in detecting LVO across diverse clinical conditions.

2. Methods

2.1. Study design and search strategy

This systematic review and meta-analysis aimed to evaluate the effectiveness of the RACE tool in detecting LVO among suspected or confirmed stroke patients. The study was guided by the PICO (patient, intervention, comparison, outcome) framework, where P referred to suspected or confirmed stroke patients, I to the RACE scoring tool, C to radiographic imaging techniques such as CT angiography (CTA), Magnetic Resonance Imaging (MRI), Transcranial Doppler ultrasound (TCD) or Digital Subtraction Angiography (DSA), and O to patients with final diagnosis of LVO. A comprehensive search strategy was employed, covering databases including Medline (via PubMed), Scopus, Embase, and Web of Science, up until June 25th, 2023. Relevant keywords related to "RACE" and "Stroke" were selected using MeSH and Emtree terms, guided by consultations with field experts and reviews of pertinent literature.

Additionally, manual searches were performed on the Google and Google Scholar search engines. The complete search strategy is provided in Supplementary Material 1.

2.2. Selection criteria

Inclusion criteria encompassed articles investigating the utility of the RACE tool in detecting LVO among suspected or confirmed stroke patients, irrespective of LVO definition and setting. Exclusion criteria comprised reviews, studies lacking non-LVO control groups, studies without reported cut-off points, studies with no required data, and duplicate publica-

tions.

2.3. Data collection

After eliminating duplicates, two reviewers independently assessed the retrieved records. The screening process was conducted in two phases: an initial evaluation of titles and abstracts, followed by a comprehensive examination of full-text articles to determine their eligibility based on selection criteria. Data from selected articles were synthesized and extracted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (16). The extracted information included study characteristics such as the first author's surname, publication year, study design, age and sex distribution of patients, sample size, the number of patients with LVO, the LVO definition used in the study, and the reference standard test employed. Additionally, diagnostic performance metrics of the RACE tool, including true positives (TP), false positives (FP), true negatives (TN), and false negatives (FN), were extracted. These metrics were subsequently used to calculate sensitivity, specificity, and other relevant variables.

2.4. Quality assessment and certainty of evidence

The Quality Assessment of Diagnostic Accuracy Studies-2 (QUADAS-2) tool (17) was used to evaluate the quality and risk of bias in the included studies. QUADAS-2 assesses bias and applicability across domains related to patient selection, index test, reference standard, and flow and timing, and each domain's risk is rated as high, unclear, or low. The certainty of evidence was determined utilizing the Grades of Recommendation, Assessment, Development, and Evaluation (GRADE) approach (18).

2.5. Statistical analysis

Statistical analyses were conducted using STATA v17.0 statistical software, with calculations performed using the 'midas' package. We used the collected TP, TN, FP, and FN values to calculate diagnostic accuracy metrics for the RACE tool, including sensitivity, specificity, the area under the receiver operating characteristic curve (AUC), diagnostic odds ratio (DOR), positive likelihood ratio (PLR), and negative likelihood ratio (NLR). Since the diagnostic value of RACE was reported at various cut-off points, we stratified the results based on reported cut-off points. In articles with multiple reports for various LVO definitions (anterior LVO and both anterior and posterior LVO) and population (suspected or confirmed stroke), the report with the broadest definition was included in the main analysis. Heterogeneity among the studies was assessed using the I² test, and we applied a bivariate random-effects model for our analysis.

Additionally, subgroup analysis was performed on studies based on criteria such as the study population (suspected or

confirmed stroke), study design, the setting in which they were conducted, assessor specialties, and the definition of LVO. Publication bias was investigated using Deeks' funnel plot asymmetry test.

3. Results

3.1. Study characteristics

A total of 345 articles were initially collected; 332 were identified through a systematic search and 13 through manual search. After duplicate removal and review of titles and abstracts, 60 articles underwent a comprehensive full-detail assessment. Ultimately, 43 articles were included in this systematic review and meta-analysis (7, 8, 11, 13, 19-57) (Figure 1). Among the selected articles, 20 were conducted in the USA, 7 in Spain, 4 in China, and 3 in the Netherlands. The rest of the articles were conducted in: Switzerland, Finland, Brazil, France, Hungary, Australia, and Denmark. One of the included articles involved multinational study populations. Twenty-one studies had a prospective design, and 21 were retrospective studies. One study had separate retrospective and prospective reports.

In total, the collective sample size encompassed 37,811 patients. The average age of participants ranged from 62.04 to 73.5 years. In 13 articles, sex-specific patient numbers were not reported. However, among the remaining articles, out of a total of 31508 patients, 16016 (50.83%) were male. Regarding the RACE cut-off points, seven articles used a cut-off point of ≥ 1 , eight used ≥ 2 , eleven used ≥ 3 , twelve used ≥ 4 , thirty-seven used ≥ 5 , ten used ≥ 6 , nine used ≥ 7 , eight used ≥ 8 , and seven used ≥ 9 . RACE evaluations were conducted in a prehospital setting in 24 studies, while they were conducted in an in-hospital setting in 17 studies. One study did not specify the setting, and in another study, data were collected from both prehospital and in-hospital medical records. RACE assessments were carried out by physicians and neurologists in 12 articles and by other Emergency Medical Services (EMS) and Emergency Department (ED) staff in 25 studies. Four studies did not mention the assessors, and in two studies, it was noted that the assessors were members of the stroke team certified in National Institutes of Health Stroke Scale (NIHSS) or were certified research personnel, although their specific specialty remained unclear. In 20 studies both anterior and posterior circulations were evaluated for LVO, whereas the definition of LVO was restricted to anterior circulation occlusions in 7 studies. Two studies, evaluated their study population based on each of the above definitions of LVO, separately. Additionally, in 14 studies, the exact definition of LVO was not provided. Table 1 provides the characteristics of the included studies.

3.2. Meta-analysis

3.2.1. Diagnostic value of RACE in detection of LVO

In the meta-analysis of RACE for LVO detection, performance varied across different cut-off points. Due to scarcity of data, conducting an analysis at ≥ 1 cut-off was not possible. For ≥ 2 cut-off, RACE had an AUC of 0.71, sensitivity of 0.94, and specificity of 0.33 with a DOR of 7. The ≥ 3 cut-off yielded an AUC of 0.79, sensitivity of 0.86, specificity of 0.57, and DOR of 8. At ≥ 4 cut-off, the AUC was 0.79, sensitivity and specificity were 0.78 and 0.68, respectively, and DOR was 7. Raising the threshold to ≥ 5 resulted in an AUC of 0.80, sensitivity of 0.69, specificity of 0.80, and DOR of 9. The ≥ 6 cut-off had an AUC of 0.77, sensitivity of 0.6, specificity of 0.80, and DOR of 6. Using ≥ 7 cut-off, the AUC was 0.70, sensitivity was 0.43, specificity was 0.89, and DOR was 6. At ≥ 8 cut-off, sensitivity was 0.26, specificity was 0.95, and DOR was 6. Finally, for ≥ 9 cut-off, sensitivity was 0.12, specificity was 0.99, and DOR was 10 (Table 2). Due to the low sensitivity observed at cut-offs ≥ 8 and ≥ 9 , the AUC values for these specific cut-off points are not reported in the results and the table (Supplementary Figure 1-9).

After evaluating the AUC, sensitivity, and specificity, the optimal cut-off points for the RACE scale in detecting LVO were determined to be ≥ 3 and ≥ 4 . However, since the majority of articles used a cut-off of ≥ 5 as the optimal, further analyses were conducted using this cut-off.

3.2.2. Subgroup analysis

At the cut-off point of ≥ 5 , there was significant heterogeneity among the studies ($I^2 = 100.0\%$). As a result, we performed a meta-regression analysis to explore the potential factors responsible for this observed heterogeneity across the studies. Our analysis revealed that the most likely contributors to this heterogeneity included variations in study settings, the specialties of the assessors using the RACE scale, and the definition of LVO ($p < 0.001$).

Subgroup analysis unveiled significant sensitivity and specificity variations based on settings, assessor specialty, population, LVO definition, and study designs. Higher sensitivity was observed when RACE investigated suspected stroke patients ($p < 0.001$), in pre-hospital settings ($p < 0.001$), when considering both anterior and posterior circulation occlusions in LVO definition ($p < 0.01$), and in prospective study designs ($p < 0.01$). Conversely, higher specificity was found when RACE investigated confirmed stroke patients ($p < 0.01$), in in-hospital settings ($p < 0.01$), only anterior circulation occlusions was considered in LVO definition ($p < 0.01$), and in retrospective study designs ($p < 0.001$). Furthermore, according to the analysis, both sensitivity ($p < 0.001$) and specificity ($p < 0.001$) were higher when the assessor was a neurologist or physician compared to other ED and EMS staff. Despite these variations, our study found comparable diagnostic ac-

curacy across different conditions (Table 3).

3.3. Publication bias

The Deeks' funnel asymmetry test demonstrated no evidence of publication bias or asymmetry in cut-off points ≥ 2 ($p=0.6$), ≥ 3 ($p=0.31$), ≥ 4 ($p=0.74$), ≥ 5 ($p=0.2$), ≥ 6 ($p=0.08$), ≥ 7 ($p=0.81$), ≥ 8 ($p=0.96$), ≥ 9 ($p=0.89$) (Supplementary Figure 10).

3.4. Risk of bias assessment

The QUADAS-2 guidelines were used to assess the quality of the included articles. In patient selection, 4 studies had high risk of bias due to inappropriate exclusions. In the Index test domain, 10 articles had unclear risk due to no mention of assessor blinding. In the reference standard section, 16 studies had unclear risk: 6 due to no mention of the utilized reference standard, 10 due to no mention of reference standard assessor blinding. In flow and timing, 11 articles had unclear risk due to uncertain patient inclusion in the final analysis (Table 4).

3.5. Certainty of evidence

The certainty of evidence in the included articles was evaluated in accordance with the GRADE guidelines. These articles were structured as observational studies, and in line with the GRADE guidelines, the initial level of evidence was set as low. The included studies exhibited significant risk of bias and substantial heterogeneity. Additionally, the presence of a very large magnitude effect size, plausible confounders and dose-response gradient effect could potentially elevate the level of evidence by four points: two for the very large magnitude effect size, one for plausibility, and one for the dose-response gradient effect. The presence of a plausible confounder is due to the enhanced diagnostic value achieved through the consideration of assessor-based grouping. Additionally, the existence of a dose-response gradient effect is a result of evaluating the performance of this scale at various cut-off points. Consequently, the certainty of evidence for the predictive value of the RACE scale in LVO detection was classified as high (Table 5).

4. Discussion

The current review assessed the RACE scale's diagnostic efficacy in detecting LVOs. Our result demonstrated that the optimal cut-off for the RACE scale is set at ≥ 3 and ≥ 4 , with a sensitivity of 0.86 (95% CI: 0.78, 0.91) and specificity of 0.57 (95% CI: 0.49, 0.67) for cut-off 3, and a sensitivity of 0.78 (95% CI: 0.70, 0.84) and specificity of 0.68 (95% CI: 0.59, 0.75) for cut-off 4.

In 2014, de la Ossa and colleagues introduced the RACE scale for the first time, building upon the components of the

NIHSS. The primary goal of developing this tool was to provide paramedics with an easy to use and accurate method of identifying patients with LVOs and facilitating the rapid transfer to CSCs. The RACE scale assigns scores based on assessments of facial palsy (scored from 0 to 2), arm motor function (scored from 0 to 2), leg motor function (scored from 0 to 2), gaze (scored from 0 to 1), and aphasia or agnosia (scored from 0 to 2). In their study, the authors sought to validate the RACE scale in a prehospital setting, where it was administered by trained medical emergency technicians as part of a prospective study. The results revealed an AUC of 0.82, a sensitivity of 0.85, and a specificity of 0.68 (7). Subsequently, numerous studies have been conducted with varying study designs, in different clinical settings, and involving various clinical specialties to validate its utility in diverse situations. A review conducted by Smith et al. assessed the performance of various scales in identifying LVO in both suspected and confirmed stroke patients. They found that, based on the analysis of three studies, the RACE score of ≥ 5 demonstrated a sensitivity and specificity of 0.69 and 0.81, respectively, for suspected stroke patients. Additionally, in confirmed stroke patients, the review revealed a sensitivity of 0.67 and a specificity of 0.85 based on an analysis of two articles. It's important to note that this review had limitations due to the relatively small number of included articles. In contrast, our review provides more robust results as it incorporates a larger number of articles for analysis (15).

In de la Ossa's study, the RACE scale was originally developed and validated in a prehospital setting, carried out by EMS technicians for individuals with suspected stroke (7). However, our subgroup analysis in this study revealed variations in sensitivity and specificity across different conditions. Nevertheless, the RACE scale demonstrated relatively comparable diagnostic accuracy across these variations. The results of our study indicated that when administered by neurologists and ED physicians, the sensitivity and specificity of the RACE scale is slightly superior compared to its use by other ED or EMS staff. This difference could be attributed to the higher level of knowledge and expertise possessed by neurologists in assessing stroke patients, as compared to other healthcare providers in a clinical setting. This could underscore the importance of training the ED and EMS staff for using RACE scale. Our study also revealed that RACE exhibits higher specificity when conducted in confirmed AIS patients. This supports previous findings that the moderate specificity of the RACE scale may result from including patients with hemorrhagic stroke, who typically exhibit severe symptoms and high RACE scores (7).

The de la Ossa study identified a cut-off of ≥ 5 as the optimal for identifying LVO patients using the RACE scale (7). As a result, many subsequent studies have focused on evaluating the diagnostic performance of RACE specifically at this

cut-off. However, our study prioritized not missing LVO patients, considering the severe consequences in terms of mortality and morbidity if the therapeutic window is missed. To address this, we recommend utilizing cut-offs of ≥ 3 or ≥ 4 , which showed higher sensitivity rates at the expense of lower specificity rates. This approach aims to ensure that LVO patients are directed to CSCs where they can receive appropriate care. It is crucial to note that some patients classified as false positives are actually hemorrhagic stroke patients, who also benefit from referral to CSCs. Therefore, using cut-offs of ≥ 3 and ≥ 4 has the potential to improve outcomes for suspected patients.

The ideal prediction tool for LVOs should incorporate several key features to ensure its effectiveness in a prehospital setting. First and foremost, the tool should prioritize simplicity and clarity to reduce the risk of misinterpretation when responding to different commands. It is crucial to employ a binary scoring system rather than relying on ordinal gradations. Speed is paramount to expedite decision-making and minimize time delays. Furthermore, the tool's design should ensure reproducibility among different healthcare personnel, which can be achieved by excluding items that might pose scoring challenges. In this regard, nuanced findings such as neglect should be avoided to ensure easy usability. RACE scale is a short and fast screening tool containing items that examine gaze deviation and palsy and motor symptoms that are the most predictive subitems of LVO and it was validated in a prospective prehospital validation cohort (1, 58-60).

The RACE test has certain limitations, notably it's a non-binary scoring system with three possible responses for most items, complicating the distinction between mild, moderate, and severe deficits and requiring nuanced interpretation. Additionally, the inclusion of assessments for aphasia and agnosia/neglect in the RACE test has been shown to demand the expertise of trained and experienced staff for accurate evaluation, potentially limiting its practicality due to the need for specialized skills and the relatively modest gains in diagnostic accuracy.

The de la Ossa study underscores the strong correlation between RACE and NIHSS, but it's crucial to note that RACE's distinct scoring and focus may not perform as effectively as NIHSS in LVO detection. In this regard, it's important to note that RACE places greater emphasis on motor symptoms, allowing for an additional point assessment for facial weakness and up to two extra points for leg weakness. However, motor symptoms, while associated with higher NIHSS scores, may not effectively differentiate between non-LVOs and LVOs, as they can also manifest in subcortical or lacunar strokes. Conversely, the RACE scale assigns only one point for gaze deviation, a characteristic sign of cortical or brain stem dysfunction and a potent LVO stroke discriminator. Further-

more, while NIHSS evaluates both fluency and comprehension, RACE focuses solely on speech with commands, potentially missing expressive aphasia—a strong LVO stroke indicator. Lastly, the RACE scale limits aphasia evaluation to individuals with right weakness and neglect to those with left-sided weakness, overlooking concurrent neglect and aphasia possibilities and left-handed patients with right hemisphere dominance (13).

While our findings provide valuable insights, this study has some limitations. In our study, performing a meta-analysis for a cut-off of ≥ 1 was not feasible. Furthermore, the role of staff training was inadequately addressed in the majority of the included studies and limited us in evaluating the role of training in diagnostic accuracy of RACE. Additionally, a notable number of studies exhibited a serious risk of bias, raising concerns about the overall quality of the evidence.

5. Conclusion

A high level of evidence indicates that the RACE scale lacks promising diagnostic value for detection of LVOs. A sensitivity range of 0.69-0.86 is insufficient for a screening tool intended to aid in the diagnosis of strokes, considering the substantial morbidity and mortality associated with this condition.

6. Declarations

6.1. Acknowledgments

The completion of this article would not have been possible without the valuable contributions and expertise of Dr. Mahmoud Yousefifard and Dr. Koohyar Ahmadzadeh. We appreciate their collaboration and guidance throughout the research process.

6.2. Conflict of interest

The authors declare no conflict of interest.

6.3. Funding and supports

None.

6.4. Authors' contribution

Both authors had similar contributions in all steps of the study. The two authors have read and approved the final version of manuscript.

6.5. Using artificial intelligence chatbots

None.

References

- Nicholls JK, Ince J, Minhas JS, Chung EML. Emerging Detection Techniques for Large Vessel Occlusion Stroke: A Scoping Review. *Front Neurol*. 2021;12:780324.
- Malhotra K, Gornbein J, Saver JL. Ischemic Strokes Due to Large-Vessel Occlusions Contribute Disproportionately to Stroke-Related Dependence and Death: A Review. *Front Neurol*. 2017;8:651.
- Nogueira RG, Jadhav AP, Haussen DC, Bonafe A, Budzik RF, Bhuva P, et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N Engl J Med*. 2018;378(1):11-21.
- Powers WJ, Derdeyn CP, Biller J, Coffey CS, Hoh BL, Jauch EC, et al. 2015 American Heart Association/American Stroke Association Focused Update of the 2013 Guidelines for the Early Management of Patients With Acute Ischemic Stroke Regarding Endovascular Treatment: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2015;46(10):3020-35.
- Albers GW, Marks MP, Kemp S, Christensen S, Tsai JP, Ortega-Gutierrez S, et al. Thrombectomy for Stroke at 6 to 16 Hours with Selection by Perfusion Imaging. *N Engl J Med*. 2018;378(8):708-18.
- Southerland AM, Johnston KC, Molina CA, Selim MH, Kamal N, Goyal M. Suspected Large Vessel Occlusion: Should Emergency Medical Services Transport to the Nearest Primary Stroke Center or Bypass to a Comprehensive Stroke Center With Endovascular Capabilities? *Stroke*. 2016;47(7):1965-7.
- Pérez de la Ossa N, Carrera D, Gorchs M, Querol M, Millán M, Gomis M, et al. Design and validation of a prehospital stroke scale to predict large arterial occlusion: the rapid arterial occlusion evaluation scale. *Stroke*. 2014;45(1):87-91.
- Noorian AR, Sanossian N, Shkirkova K, Liebeskind DS, Eckstein M, Stratton SJ, et al. Los Angeles Motor Scale to Identify Large Vessel Occlusion: Prehospital Validation and Comparison With Other Screens. *Stroke*. 2018;49(3):565-72.
- Katz BS, McMullan JT, Sucharew H, Adeoye O, Broderick JP. Design and validation of a prehospital scale to predict stroke severity: Cincinnati Prehospital Stroke Severity Scale. *Stroke*. 2015;46(6):1508-12.
- Venizelos A, Chen SH, Gianatasio R, Coffman SR, Gamber MA, Myers JH, Janardhan V. Abstract T MP59: Texas Stroke Intervention Pre-Hospital Stroke Severity Scale (aka LEGS score): A Triaging Tool for Interventional Stroke Therapy. *Stroke*. 2014;45.
- Hastrup S, Damgaard D, Johnsen SP, Andersen G. Prehospital Acute Stroke Severity Scale to Predict Large Artery Occlusion: Design and Comparison With Other Scales. *Stroke*. 2016;47(7):1772-6.
- Teleb MS, Ver Hage A, Carter J, Jayaraman MV, McTaggart RA. Stroke vision, aphasia, neglect (VAN) assessment-a novel emergent large vessel occlusion screening tool: pilot study and comparison with current clinical severity indices. *J Neurointerv Surg*. 2017;9(2):122-6.
- Lima FO, Silva GS, Furie KL, Frankel MR, Lev MH, Caramargo É C, et al. Field Assessment Stroke Triage for Emergency Destination: A Simple and Accurate Prehospital Scale to Detect Large Vessel Occlusion Strokes. *Stroke*. 2016;47(8):1997-2002.
- Heldner MR, Zubler C, Mattle HP, Schroth G, Weck A, Mono ML, et al. National Institutes of Health stroke scale score and vessel occlusion in 2152 patients with acute ischemic stroke. *Stroke*. 2013;44(4):1153-7.
- Smith EE, Kent DM, Bulsara KR, Leung LY, Lichtman JH, Reeves MJ, et al. Accuracy of Prediction Instruments for Diagnosing Large Vessel Occlusion in Individuals With Suspected Stroke: A Systematic Review for the 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke*. 2018;49(3):e111-e22.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg*. 2010;8(5):336-41.
- Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*. 2011;155(8):529-36.
- Yang B, Mustafa RA, Bossuyt PM, Brozek J, Hultcrantz M, Leeflang MMG, et al. GRADE Guidance: 31. Assessing the certainty across a body of evidence for comparative test accuracy. *J Clin Epidemiol*. 2021;136:146-56.
- Zhang B, Huo X, Yuan F, Song G, Liu L, Ma G, et al. Design and validation of a recognition instrument-the stroke aid for emergency scale-to predict large vessel occlusion stroke. *Aging (Albany NY)*. 2021;13(10):13680-92.
- Zaidi SF, Shawver J, Espinosa Morales A, Salahuddin H, Tietjen G, Lindstrom D, et al. Stroke care: initial data from a county-based bypass protocol for patients with acute stroke. *J Neurointerv Surg*. 2017;9(7):631-5.
- Turc G, Maier B, Naggara O, Seners P, Isabel C, Tisserand M, et al. Clinical Scales Do Not Reliably Identify Acute Ischemic Stroke Patients With Large-Artery Occlusion. *Stroke*. 2016;47(6):1466-72.
- Thavarajah S, Langston Z, Sarayusa A, Fowler LA, Sivakumar S, Shah N. Evaluation of the Rapid Arterial Occlusion Evaluation (RACE) scale in Upstate South Carolina, USA. *J Stroke Cerebrovasc Dis*. 2022;31(11):106746.
- Tárkányi G, Karádi ZN, Csécséi P, Bosnyák E, Fehér G, Molnár T, Szapáry L. Capability of stroke scales to detect large vessel occlusion in acute ischemic stroke - a pilot

- study. *Ideggyogy Sz.* 2021;74(3-4):99-103.
24. Sequeira D, Martin-Gill C, Guyette FX. APJ. Abstract 12493: Comparison of Prehospital Stroke Scales| Circulation. *Circulation [Internet]*. 2015; 132: A12493.
 25. Scheitz JF, Abdul-Rahim AH, MacIsaac RL, Cooray C, Sucharew H, Kleindorfer D, et al. Clinical Selection Strategies to Identify Ischemic Stroke Patients With Large Anterior Vessel Occlusion: Results From SITS-ISTR (Safe Implementation of Thrombolysis in Stroke International Stroke Thrombolysis Registry). *Stroke*. 2017;48(2):290-7.
 26. Rodríguez-Pardo J, Fuentes B, Alonso de Leciana M, Ximénez-Carrillo Á, Zapata-Wainberg G, Álvarez-Fraga J, et al. The Direct Referral to Endovascular Center criteria: a proposal for pre-hospital evaluation of acute stroke in the Madrid Stroke Network. *Eur J Neurol*. 2017;24(3):509-15.
 27. Qureshi S, Janjua AU, Raza SA, Brasher C, Anderson A, Belagaje S, et al. Abstract WMP67: Stroke Screening Tools Have High Specificity for Detecting Large Vessel Occlusion in a Southeastern US Prospective Cohort Study. *Stroke*. 2016;47(suppl_1):AWMP67-AWMP.
 28. Nguyen TTM, van den Wijngaard IR, Bosch J, van Belle E, van Zwet EW, Dofferhoff-Vermeulen T, et al. Comparison of Prehospital Scales for Predicting Large Anterior Vessel Occlusion in the Ambulance Setting. *JAMA Neurol*. 2021;78(2):157-64.
 29. Navakele D, Vahidy F, Kendrick S, Traylor A, Haydel M, Drury S, Martin-Schild S. Vision, Aphasia, Neglect Assessment for Large Vessel Occlusion Stroke. *J Stroke Cerebrovasc Dis*. 2020;29(1):104478.
 30. Mehta BP, Jadhav A, Sessa J, Katz R, Duong H, Lima A, et al. Assessment of the Rapid arterial occlusion evaluation (RACE) scale in real-world practice for prediction of large vessel occlusion and reducing time to thrombectomy. *Stroke*. 2018;49(Suppl_1):A96-A.
 31. Mayasi Y, Goddeau RP, Jr., Moonis M, Silver B, Jun-O'Connell AH, Puri AS, Henninger N. Leukoaraiosis Attenuates Diagnostic Accuracy of Large-Vessel Occlusion Scales. *AJNR Am J Neuroradiol*. 2018;39(2):317-22.
 32. Keenan KJ, Smith WS, Cole SB, Martin C, Hemphill JC, Madhok DY. Large vessel occlusion prediction scales provide high negative but low positive predictive values in prehospital suspected stroke patients. *BMJ Neurol Open*. 2022;4(2):e000272.
 33. Keenan KJ, Smith WS. The Speech Arm Vision Eyes (SAVE) scale predicts large vessel occlusion stroke as well as more complicated scales. *J Neurointerv Surg*. 2019;11(7):659-63.
 34. Jumaa MA, Castonguay AC, Salahuddin H, Shawver J, Sajū L, Burgess R, et al. Long-term implementation of a prehospital severity scale for EMS triage of acute stroke: a real-world experience. *J Neurointerv Surg*. 2020;12(1):19-24.
 35. James R, Cruz A, Fortuny E, Ugiliweneza B, Wang D, White A, et al. E-063 Primary results from the predict (pre-hospital emergency IVO detection during initial care transport) study: a prospective, multi-center, non-inferiority comparison between the novel PREDICT composite 4-item scale and the rapid arterial occlusion evaluation (RACE) scale to detect large-vessel occlusion (LVO) strokes when administered by us-based ems personnel to suspected stroke patients at the initial pre-hospital encounter. *British Medical Journal Publishing Group*; 2020.
 36. Heldner MR, Hsieh K, Broeg-Morvay A, Mordasini P, Bühlmann M, Jung S, et al. Clinical prediction of large vessel occlusion in anterior circulation stroke: mission impossible? *J Neurol*. 2016;263(8):1633-40.
 37. Gropen TI, Boehme A, Martin-Schild S, Albright K, Samai A, Pishanidar S, et al. Derivation and Validation of the Emergency Medical Stroke Assessment and Comparison of Large Vessel Occlusion Scales. *J Stroke Cerebrovasc Dis*. 2018;27(3):806-15.
 38. Gong X, Chen Z, Shi F, Zhang M, Xu C, Zhang R, Lou M. Conveniently-Grasped Field Assessment Stroke Triage (CG-FAST): A Modified Scale to Detect Large Vessel Occlusion Stroke. *Front Neurol*. 2019;10:390.
 39. Globler NK, Fuls T, Supples M, Panagos P, Kim D. Factors Predicting Misidentification of Acute Ischemic Stroke and Large Vessel Occlusion by Paramedics. *Crit Pathw Cardiol*. 2022;21(4):172-5.
 40. English SW, Chhabra N, Hanus AE, Basharath R, Miller M, Butterfield RJ, et al. Prehospital stroke scales outperform National Institutes of Health Stroke Scale in predicting large vessel occlusion in a large academic telestroke network. *J Telemed Telecare*. 2023;1357633x231204066.
 41. Duvekot MHC, Venema E, Rozeman AD, Moudrou W, Vermeij FH, Biekart M, et al. Comparison of eight prehospital stroke scales to detect intracranial large-vessel occlusion in suspected stroke (PRESTO): a prospective observational study. *Lancet Neurol*. 2021;20(3):213-21.
 42. Dickson RL, Crowe RP, Patrick C, Crocker K, Aiken M, Adams A, et al. Performance of the RACE Score for the Prehospital Identification of Large Vessel Occlusion Stroke in a Suburban/Rural EMS Service. *Prehosp Emerg Care*. 2019;23(5):612-8.
 43. Dickson R, Gleisberg G, Aiken C, Crocker K, Mason C, Nichols T, Patrick C. 7 Validation and Integration of a Stroke Algorithm (VISA Study): Out-of-Hospital Identification of Large Vessel Occlusion Stroke in a Suburban/Rural EMS Service. *Annals of Emergency Medicine*. 2017;70(4):S3-S4.
 44. Perez de la Ossa N, Abilleira S, Ribó M, Monica M, Cardona P, Urra X, et al. External Validation of the RACE Scale

- After Its Implementation in the Stroke Code Protocol in Catalonia. *Stroke*. 2017;48(suppl_1):A18-A.
45. Pérez de la Ossa N, Ribó M, Jiménez X, Abilleira S. Pre-hospital Scales to Identify Patients With Large Vessel Occlusion: It Is Time for Action. *Stroke*. 2016;47(11):2877-8.
 46. Dekker L, Geraedts VJ, Hubert J, Duijndam D, Durieux MD, Janssens L, et al. Prehospital Triage of Intracranial Hemorrhage and Anterior Large-Vessel Occlusion Ischemic Stroke: Value of the Rapid Arterial Occlusion Evaluation. *Stroke: Vascular and Interventional Neurology*. 2023:e000947.
 47. Cruz A, Fortuny E, Ugiliweneza B, Wang D, White A, Khatrar N, et al. O-004 Prospective, multi-centered, EMS-administered, Pre-hospital validation study of the rapid arterial occlusion evaluation (RACE) scale for detecting large vessel occlusion stroke in the United States compared to the original RACE validation study from Spain: a subanalysis of the PREDICT study. *British Medical Journal Publishing Group*; 2020.
 48. Crowe RP, Myers JB, Fernandez AR, Bourn S, McMullan JT. The Cincinnati Prehospital Stroke Scale Compared to Stroke Severity Tools for Large Vessel Occlusion Stroke Prediction. *Prehosp Emerg Care*. 2021;25(1):67-75.
 49. Chiu YC, Hsieh MJ, Lin YH, Tang SC, Sun JT, Chiang WC, et al. External validation of prehospital stroke scales for emergent large vessel occlusion. *Am J Emerg Med*. 2021;41:35-9.
 50. Chen Z, Zhang R, Xu F, Gong X, Shi F, Zhang M, Lou M. Novel Prehospital Prediction Model of Large Vessel Occlusion Using Artificial Neural Network. *Front Aging Neurosci*. 2018;10:181.
 51. CHEN H, ZHAO Y, WANG C, MU Z, LIU H, HAN W. Predictive value of modified RACE score for large vessel occlusion in patients with acute anterior circulation ischemic stroke. *International Journal of Cerebrovascular Diseases*. 2020:241-6.
 52. Abstracts of the 22nd European Stroke Conference. London, United Kingdom. May 28-31, 2013. *Cerebrovasc Dis*. 2012;35 Suppl 3:9-854.
 53. Carrera D, Gorchs M, Querol M, Abilleira S, Ribó M, Millán M, et al. Revalidation of the RACE scale after its regional implementation in Catalonia: a triage tool for large vessel occlusion. *J Neurointerv Surg*. 2019;11(8):751-6.
 54. Carrera D, Campbell BC, Cortés J, Gorchs M, Querol M, Jiménez X, et al. Predictive Value of Modifications of the Prehospital Rapid Arterial Occlusion Evaluation Scale for Large Vessel Occlusion in Patients with Acute Stroke. *J Stroke Cerebrovasc Dis*. 2017;26(1):74-7.
 55. Anadani M, Almallouhi E, Wahlquist AE, Debenham E, Holmstedt CA. The Accuracy of Large Vessel Occlusion Recognition Scales in Telestroke Setting. *Telemed J E Health*. 2019;25(11):1071-6.
 56. Zhao H, Coote S, Pesavento L, Churilov L, Dewey HM, Davis SM, Campbell BC. Large Vessel Occlusion Scales Increase Delivery to Endovascular Centers Without Excessive Harm From Misclassifications. *Stroke*. 2017;48(3):568-73.
 57. Puolakka T, Virtanen P, Kuusma M, Strbian D. Comparison of large vessel occlusion scales using prehospital patient reports. *Acta Neurol Scand*. 2022;145(3):265-72.
 58. Keenan KJ, Kircher C, McMullan JT. Prehospital Prediction of Large Vessel Occlusion in Suspected Stroke Patients. *Curr Atheroscler Rep*. 2018;20(7):34.
 59. El-Ghanem M, Al-Mufti F, Thulasi V, Singh IP, Gandhi C. Expanding the treatment window for ischemic stroke through the application of novel system-based technology. *Neurosurg Focus*. 2017;42(4):E7.
 60. van Gaal S, Demchuk A. Clinical and Technological Approaches to the Prehospital Diagnosis of Large Vessel Occlusion. *Stroke*. 2018;49(4):1036-43.







Figure 1: Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of current meta-analysis; LVO: Large vessels occlusion.




Supplementary figure 1: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 2 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.




Supplementary figure 2: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 3 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.




Supplementary figure 3: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 4 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.




Supplementary figure 4: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 5 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.




Supplementary figure 5: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 6 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.









Supplementary figure 6: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 7 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.











Supplementary figure 7: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 8 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.



Supplementary figure 8: Sensitivity, specificity, and diagnostic odds ratio of RACE in cut-off point of ≥ 9 , in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; CI: confidence interval.

cut-off ≥ 2 **cut-off ≥ 3** **cut-off ≥ 4** **cut-off ≥ 5** **cut-off ≥ 6** **cut-off ≥ 7** 

Supplementary figure 9: SROC and area under the curve of RACE in different cut-off points, in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion; SROC: summary receiver operating characteristic.

cut-off ≥ 2 **cut-off ≥ 3** **cut-off ≥ 4** **cut-off ≥ 5** **cut-off ≥ 6** **cut-off ≥ 7** **cut-off ≥ 8** **cut-off ≥ 9** 

Supplementary figure 10: Publication bias of RACE in different cut-off points, in detecting LVO. RACE: Rapid Arterial Occlusion Evaluation; LVO: Large vessel occlusion.

Table 1: Characteristics of the included studies

Author	Setting	Assessor	Design	Population	Cut-off	Sample size	Male (n)	Mean age	References	IVO definition
Anadani, 2019, USA	Prehospital	Neurologists	Retrospective	SS	5	439	231	66.7	CTA	ICA, MCA (M1, M2), PCA (P1), ACA (A1), VA, BA
Carrera, 2017, Spain	Prehospital	EMS staff	Prospective	SS	5	341	182	70	CTA, MRA or arteriography	ICA, MCA, BA
Carrera, 2018, Spain	Prehospital	EMS staff	Prospective	SS	1-9	1822	957	73.5	CTA, MRA or TSD	ICA, MCA (M1), BA
Carrera, 2013, Spain	Prehospital	Prehospital personnel	Retrospective	SS	4	263	NR	NR	TSD or cerebral angiography	NR
Chen, 2020, China	NR	NR	Prospective	CS	5	184	NR	62.04	CT	NR
Chen, 2018, China	ED	Experienced doctors	Retrospective	CS	4	600	368	67.5	CTA or TOF-MRA	ICA, MCA (M1, M2), BA
Chiu, 2021, Taiwan	ED	Nurses	Retrospective	CS	5	1231	704	70.5	CTA or MRA	ICA, MCA (M1, M2) ACA (A1), PCA (P1), BA
Crowe, 2020, USA	Prehospital	EMS staff	Retrospective	SS	1-9	1949	903	NR	CTA or MRI	ICA, MCA, BA
Cruz, 2020, USA	Prehospital	EMS personnel	Prospective	SS	6,7,8	232	NR	NR	CTA, MRA or DSA	NR
Dekker, 2023, Netherlands	Prehospital	Ambulance paramedics and registered nurses with specialized training in ambulance care	Prospective	SS	1-9	2004	1020	71.1	NR	ICA, MCA (M1, M2), ACA (A1, A2)
De la Ossa, 2014, Spain	Prehospital	Trained medical emergency technicians	Prospective	SS	1-9	357	193	73	CTA, MRA or TSD	ICA, MCA (M1), BA
De la Ossa, 2016, Spain	Prehospital	EMS staff	Prospective	SS	5	749	NR	NR	NR	NR
De la Ossa, 2017, Spain	Prehospital	EMS staff	Prospective	SS	5	962	NR	NR	NR	ICA, MCA (M1, M2), BA
Dickson, 2017, USA	Prehospital	EMS staff	Prospective	SS	5	161	NR	NR	CTA	NR
Dickson, 2019, USA	Prehospital	EMS professionals	Retrospective	SS	1-9	440	214	NR	NR	NR
Duvekot, 2021, Netherlands	Prehospital	Ambulance paramedics trained nurses	Prospective	SS	5	1039	560	NR	CT, CTA, CT perfusion, MRI, or DSA	ICA, MCA (M1, M2), ACA (A1, A2)
English, 2023, USA	Prehospital	Neurologists	Retrospective	SS	5	625	323	67.9	CTA or MRA	ICA, MCA (M1, M2), ACA (A1, A2) ICA, MCA (M1, M2), ACA (A1, A2), PCA (P1, P2), BA

Table 1: Characteristics of the included studies (continue)

Author	Setting	Assessor	Design	Population	Cut-off	Sample size	Male (n)	Mean age	References	IVO definition
Glober, 2022, USA	In-hospital and Pre-hospital	Paramedics	Retrospective	SS	5	226	98	NR	CTA	NR
Gong, 2019, China	Hospital admission	Neurologists	Retrospective	CS	5	1355	821	69	CTA or TOF-MRA	ICA, MCA (M1, M2), BA
Gropen, 2018, USA	Stroke Center	Experienced paramedics	Retrospective	CS	5	1663	898	62	CTA or MRA	ICA, MCA, BA
Hastrup, 2016, Denmark	Hospital admission	NR	Retrospective	CS	5	3127	1868	69	CTA or MRA	ICA, MCA (M1, M2), ACA (A1, A2), PCA (P1, P2), BA
Heldner, 2016, Switzerland	Stroke Center	Neurologists	NR	CS	3	1085	658	67.7	CTA or MRA	ICA, MCA (M1, M2, M3, M4), ACA
James, 2020, USA	Prehospital	NR	Prospective	SS	5,6	232	NR	NR	CTA, MRA or DSA	ICA, MCA (M1, M2), BA
Jumaa, 2020, USA	Prehospital	EMS personnel	Prospective	SS	5	2635	219		CTA	NR
Keenan, 2019, USA	Hospital admission	Neurologists and emergency physicians	Retrospective	SS	1-9	735	NR	NR	CTA	ICA, MCA (M1, M2), BA
Keenan, 2022, USA	ED	ED nurses	Retrospective	SS	3,5,7	184	94	NR	CT or CTA	ICA, MCA (M1, M2), BA
Lima, 2016, Brazil	Hospital admission	Certified research personnel	Prospective	SS	5	727	378	68.1	CT or CTA	ICA, MCA (M1, M2), BA
Mayasi, 2017, USA	Hospital admission	Members of the stroke team certified in the NIHSS	Retrospective	CS	5	274	148	69	CTA or MRA	ICA, MCA(M1)
Mehta, 2018, USA	ED	EMS staff	Prospective	SS	5	797	NR	NR	CT	NR
Navalkel, 2019, USA	Prehospital	ED nurses	Retrospective	CS	5	224	99	66	CTA or MRA	ICA, MCA, ACA, PCA, VA, BA
Nguyen, 2020, Netherlands	Prehospital	EMS paramedics	Prospective	SS	5	2007	1021	71.1	NR	ICA, MCA (M1, M2), ACA (A1, A2)
Noorian, 2018, USA	Prehospital	Ambulance paramedics	Prospective	SS	5	94	NR	70	CTA or MRA	ICA, MCA (M1, M2), ACA (A1), PCA (P1), VA, BA

Table 1: Characteristics of the included studies (continue)

Author	Setting	Assessor	Design	Population	Cut-off	Sample size	Male (n)	Mean age	References	LVO definition
Puolakka, 2022, Finland	Prehospital	Prehospital emergency physicians and Stroke Neurologist	Retrospective	SS	5	509	NR	NR	CTA	ICA, MCA (M1, M2), BA
Qureshi, 2016, USA	Prehospital	EMS staff	Prospective	SS CS	5	92 42	NR	NR	Intracranial large artery imaging	NR
Rodriguez-Pardo, 2017, Spain(A)	Stroke Center	EMS or neurologist	Retrospective	SS	5	317	190	71.3	NR	NR
Rodriguez-Pardo, 2017, Spain(B)	Stroke Center	Neurologist	Prospective	SS	5	153	74	71.2	NR	NR
Scheitz, 2017, Multi-national	Stroke Center	Stroke physicians	Retrospective	CS	2,3,5,6	3505	1967	68.1	NR	ICA, MCA(M1), MCA (M1), BA
Sequeira, 2015, USA	Prehospital	EMS provider	Retrospective	SS	4	1293	NR	NR	NR	NR
TÁRKÁNYI, 2021, Hungary	Stroke Center	NR	Prospective	CS	4,5	180	94	68.2	CTA	ICA, MCA (M1, M2, M3), ACA (A1, A2), PCA, VA, BA
Thavarajah, 2022, USA	Prehospital	EMS personnel	Retrospective	SS	1-9	126	63	69	CTA	ICA, MCA (M1), BA ICA, MCA (M1, M2), BA
Turc, 2016, France	Hospital admission	neurologist	Retrospective	CS	5	1004	575	NR	CTA or MRA	ICA, MCA (M1, M2), BA
Zaidi, 2017, USA	Prehospital	EMS personnel	Prospective	SS	5	251	119	NR	CT and CTA	NR
Zhang, 2021, China	ED	Experienced neurologists	Prospective	CS	3	1053	689	NR	CTA or TOF-MRA	ICA, MCA (M1, M2), BA
Zhao, 2017, Australia	ED	neurology residents or fellows certified in administering the NIHSS.	Prospective	SS	5	565	288	NR	CT or CTA	ICA, MCA (M1, M2)

ACA: Anterior cerebral artery; BA: Basilar artery; CS: Confirmed stroke; CT: Computed tomography; CTA: CT angiography; DSA: Digital Subtraction Angiography; ED: Emergency department; EMS: Emergency medical service; ICA: Internal carotid artery; LVO: Large vessel occlusion; MCA: Middle cerebral artery; MRA: Magnetic resonance angiography; MRI: Magnetic resonance imaging; NR: Not reported; NIHSS: The National Institutes of Health Stroke Scale; PCA: Posterior cerebral artery; SS: suspected stroke; TCD: Transcranial Doppler ultrasound; TOF-MRA: Time-of-flight magnetic resonance angiography; VA: Vertebral artery.

Table 2: The diagnostic performance of RACE scale in different cut-off points

cut-off	Sensitivity	Specificity	PLR	NLR	DOR	AUC
≥2	0.94 (0.87-0.97)	0.33 (0.23-0.44)	1.4 (1.2-1.6)	0.19 (0.11-0.32)	7 (4-12)	0.71 (0.67-0.75)
≥3	0.86 (0.78-0.91)	0.57 (0.47-0.67)	2 (1.7-2.4)	0.25 (0.19-0.34)	8 (6-11)	0.79 (0.76-0.83)
≥4	0.78 (0.70-0.84)	0.68 (0.59-0.75)	2.4 (2-2.8)	0.33 (0.26-0.41)	7 (6-9)	0.79 (0.75-0.82)
≥5	0.69 (0.65-0.73)	0.80 (0.76-0.83)	3.4 (2.9-4.1)	0.39 (0.35-0.43)	9 (7-11)	0.80 (0.76-0.83)
≥6	0.61 (0.54-0.67)	0.80 (0.75-0.84)	3 (2.6-3.5)	0.49 (0.43-0.56)	6 (5-8)	0.77 (0.73-0.81)
≥7	0.43 (0.38-0.49)	0.89 (0.86-0.92)	4.1 (3.3-4.9)	0.63 (0.59-0.68)	6 (5-8)	0.70 (0.66-0.74)
≥8	0.26 (0.22-0.30)	0.95 (0.92-0.97)	4.9 (3.4-6.9)	0.78 (0.75-0.81)	6 (4-9)	-
≥9	0.12 (0.10-0.14)	0.99 (0.97-0.99)	9.3 (3.8-22.8)	0.89 (0.87-0.91)	10 (4-26)	-

All measures are reported with 95% confidence interval. PLR: Positive Likelihood Ratio; NLR: Negative Likelihood Ratio; DOR: Diagnostic Odds Ratio; AUC: Area Under the Curve; RACE: Rapid Arterial Occlusion Evaluation.

Table 3: Subgroup analysis in cut-off ≥5 for RACE scale

Parameter	No*	Sensitivity	Specificity	AUC	PLR	NLR	DOR
Study setting							
Prehospital	22	0.72 (0.68 -0.77)	0.77 (0.71 - 0.82)	0.81 (0.77 - 0.84)	3.1 (2.7, 3.7)	0.36 (0.31, 0.41)	9 (7, 11)
In-hospital	15	0.64 (0.58 -0.70)	0.85 (0.80 - 0.90)	0.76 (0.72 - 0.80)	4.0 (2.9, 5.5)	0.42 (0.37, 0.48)	9 (6, 14)
Assessor							
Neurologist and physicians	23	0.70 (0.65 -0.75)	0.76 (0.70 - 0.82)	0.82 (0.79 - 0.85)	4.8 (2.8, 8.1)	0.34 (0.27, 0.44)	14 (7, 26)
Other staff	10	0.71 (0.64 -0.78)	0.85 (0.79 - 0.91)	0.79 (0.75 - 0.82)	2.9 (2.5, 3.4)	0.40 (0.35, 0.45)	7 (6, 9)
Study population							
Suspected to stroke	29	0.71 (0.67 -0.75)	0.77 (0.73 - 0.82)	0.81 (0.77 - 0.84)	3.1 (2.7, 3.6)	0.37 (0.33, 0.42)	9 (7, 10)
Confirmed stroke	10	0.64 (0.57 -0.71)	0.86 (0.81 - 0.91)	0.75 (0.71 - 0.78)	4.6 (2.8, 7.6)	0.42 (0.36, 0.50)	11 (6, 21)
Study design							
Prospective	21	0.72 (0.67 -0.77)	0.77 (0.72 - 0.83)	0.80 (0.77 - 0.84)	3.2 (2.7, 3.8)	0.36 (0.32, 0.40)	9 (7, 11)
Retrospective	18	0.66 (0.60 -0.71)	0.82 (0.77 - 0.87)	0.79 (0.75 - 0.82)	3.7 (2.7, 5.1)	0.42 (0.35, 0.50)	9 (6, 14)
IVO definition (location of occlusion)							
Anterior and posterior	22	0.69 (0.64 -0.74)	0.81 (0.76 - 0.86)	0.81 (0.77 -0.84)	3.7 (2.8,4.7)	0.38 (0.33, 0.45)	10 (7,13)
Only anterior	9	0.66 (0.58 -0.74)	0.85 (0.79 - 0.91)	0.79 (0.75 -0.82)	4.5 (3.4, 6.0)	0.40 (0.36, 0.44)	11 (8, 5)

*Number of articles included in meta-analyses. All measures are reported with 95% confidence interval. PLR: Positive Likelihood Ratio; NLR: Negative Likelihood Ratio; DOR: Diagnostic Odds Ratio; AUC: Area Under the Curve; IVO: Large Vessels Occlusion; RACE: Rapid Arterial Occlusion Evaluation.

Table 4: Risk of bias assessment

Study	Risk of bias				Applicability		
	Patient selection	Index test	Reference standard	Flow and timing	Patient selection	Index test	Reference standard
Anadani, 2019	H	U	L	L	L	L	L
Carrera, 2017	L	L	U	L	L	L	L
Carrera, 2018	L	L	U	L	L	L	L
Carrera, 2013	L	L	L	U	L	L	L
Chen, 2020	L	L	U	L	L	L	L
Chen, 2018	L	L	L	U	L	L	L
Chiu, 2021	L	L	L	L	L	L	L
Crowe, 2020	L	L	L	L	L	L	L
Cruz, 2020	L	L	L	L	L	L	L
Dekker, 2023	L	U	U	L	L	L	L
De la Ossa, 2014	L	L	L	L	L	L	L
De la Ossa, 2016	L	L	U	U	L	L	L
De la Ossa, 2017	L	L	U	U	L	L	L
Dickson, 2017	U	L	U	U	L	L	L
Dickson, 2019	L	L	U	L	L	L	L
Duvekot, 2021	L	L	L	L	L	L	L
English, 2023	L	L	L	L	L	L	L
Glober, 2022	L	L	U	L	L	L	L
Gropen, 2018	L	U	L	L	L	L	L
Gong, 2019	L	L	L	L	L	L	L
Hastrup, 2016	L	L	L	H	L	L	L
Heldner, 2016	L	L	L	L	L	L	L
James, 2020	U	L	L	U	L	L	L
Jumaa, 2020	L	L	U	L	L	L	L
Keenan, 2019	H	L	L	L	L	L	L
Keenan, 2022	L	L	L	L	L	L	L
Lima, 2016	H	L	L	L	L	L	L
Mayasi, 2017	L	L	L	L	L	L	L
Mehta, 2018	L	L	H	U	L	L	L
Navalkel, 2019	L	U	L	L	L	L	L
Nguyen, 2020	L	U	L	L	L	L	L
Noorian, 2018	L	L	L	L	L	L	L
Puolakka, 2022	L	L	L	L	L	L	L
Qureshi, 2016	L	L	U	U	L	L	L
Rodriguez-Pardo, 2017	H	L	L	L	L	L	L
Scheitz, 2017	L	U	U	L	L	L	L
Sequeira, 2015	U	U	L	U	L	L	L
TÁRKÁNYI, 2021	L	L	L	L	L	L	L
Thavarajah, 2022	L	U	L	L	L	L	L
Turc, 2016	L	L	L	L	L	L	L
Zaidi, 2017	L	U	U	U	L	L	L
Zhang, 2021	L	U	L	L	L	L	L
Zhao, 2017	U	L	U	U	L	L	L

L: Low risk, U: Unclear risk, H: High risk.

Table 5: Risk of bias assessment

Outcome	Sample size	Risk of bias	Heterogeneity (I2 value)	Indirectness	Imprecision	Publication bias	Other considerations
LVO detection	37811 Event rate (%) (23.44)	Serious	Serious	No serious	No serious	Not present	Large magnitude of effect Plausible confounders Dose-response gradient effect

LVO: Large Vessels Occlusion; RACE: Rapid Arterial Occlusion Evaluation.