

Assessment of Phenolic Contents and Antioxidant and Antibacterial Activities of Extracts from Four Varieties of Iranian Date Palm (*Phoenix dactylifera* L.) Seeds

Ramin Radfar¹, Mehdi Farhoodi¹, Ismail Ghasemi², Amin Mousavi Khaneghah^{3,4}, Farzaneh Shahrzad¹, Hedayat Hosseini^{1*}

1-National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2-Department of Plastics, Iran Polymer and Petrochemical Institute (IPPI), Tehran, Iran.

3-Department of Food Science, Faculty of Food Engineering, University of Campinas, Campinas, SP, Brazil.

4-Department of General and Applied Mathematics, Azerbaijan State Oil and Industry University, Baku, Azerbaijan.

Abstract

Background and objective: Every year, large quantities of date palm seeds are produced as byproducts in date processing and packaging industries, which is discarded or used as low-value materials for animal feeds and composts. However, these bioresources may include potentials to produce high-value added products in food industries. The major aim of the current study was to assess phenolic profiles and contents and antioxidant and antibacterial activities of four Iranian date palm seed extracts, namely Zahedi, Kabkab, Mazafati and Rabbi.

Material and methods: Total phenolic contents, phenolic compounds profile and antioxidant and antibacterial activities of extracts from four Iranian date palm seeds were assessed using Folin-Ciocalteu, reversed-phase high-performance liquid chromatography, 2, 2-diphenyl-1-picrylhydrazyl radical scavenging, agar disc diffusion and broth microdilution methods, respectively.

Results and conclusion: Total phenolic contents varied 1480-3380 mg GAE 100 g⁻¹ dw. cinnamic, chlorogenic, caffeic and 3, 5-dihydroxybenzoic acid included the primary phenolic compounds, respectively. Of the varieties, Kabkab and Mazafati seed extracts with IC₅₀ values of 16.56 and 22.6 µg ml⁻¹ demonstrated the highest and lowest radical scavenging activity, respectively. Results obtained from disc diffusion method revealed that all extracts included inhibitory effects against *Staphylococcus aureus*, but not against *Escherichia coli*. Minimum inhibitory concentration and minimum bactericidal concentration of the extracts ranged 1.56-3.125 and 3.125-12.5 mg ml⁻¹ for *Staphylococcus aureus*, respectively. Based on the findings, Iranian date seeds are good sources of extractable phenolic compounds with notable antioxidant activities, which can be used as natural additives in formulations of various products such as functional foods and dietary supplements. Furthermore, these seeds can be converted to value added products through biotechnological processes.

Conflict of interest: The authors declare no conflict of interest.

Article Information

Article history:

Received 15 Nov 2018
Revised 24 Dec 2018
Accepted 07 June 2019

Keywords:

- Antibacterial properties
- Antioxidant activity
- Date palm seed
- Phenolic profile

*Corresponding author:

Hedayat Hosseini,

Department of Food Science and Technology, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Tel: + 98-21-22376426

Fax: +98- 21-22360660

E-mail: hedayat@sbmu.ac.ir

1. Introduction

It is suggested that the date palm (*Phoenix dactylifera* L.) is one of the earliest planted trees in world [1,2]. It is cultivated in most tropical and subtropical regions worldwide, including Middle East, North Africa, the United States and Southern Europe [3]. One of these regional countries, Iran, produces more than 1 million tons of dates annually; ranked highest after Egypt, Algeria,

Saudi Arabia and Iraq [4]. The edible part of date fruit is pericarp surrounding the seed. Date seed, as one of the major byproducts produced during processing operations, constitutes approximately 10-15% of the total fruit mass [5]. Currently, date seeds are discarded or used in limited scales in animal feeds and natural fertilizers. However, there are possibilities of producing high-value added

products from these byproducts [6]. Date palm seeds (mainly in the form of hydrolysates) have been used as solid substrates for the production of various fermented products, including antibiotics (such as nisin and oxytetracycline), single cell proteins, citric acids and biofuels (from seed lipids) [7-11]. Most importantly, it is possible to produce valuable functional foods from date seed extracts [12]. These products include environmental and economic advantages for the farmers and food industries. Based on the phytochemical analysis, date seeds contain natural compounds such as carbohydrates, oils, dietary fibers, minerals and vitamins [13-14]. Moreover, biologically active compounds such as phenolic acids, anthocyanins, carotenoids, sterols and flavonoids are other bioactive components of the date seeds, which offer health improvement properties such as anti-carcinogenic [15], antioxidant [16], anti-mutagenicity [17], antibacterial and anti-inflammatory effects [18-19]. Use of natural bioactive compounds extracted from plant sources is rising in food products, mainly due to the public concerns on adverse effects of artificial additives and preservatives on human health [20]. Although studies have been carried out on date palm seed properties, there are no recently published studies on compositions and characteristics of the Iranian date seeds. The major aim of this study was to assess and compare total phenolic contents and antioxidant and antibacterial activities of four Iranian date seed varieties, namely Zahedi, Kabkab, Mazafati and Rabbi.

2. Materials and methods

2.1. Materials and bacterial strains

Folin-Ciocalteu phenol reagent, 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH), gallic acid, cinnamic acid, caffeic acid, ferulic acid, vanillic acid, dihydroxybenzoic acids and chlorogenic acid were purchased from Sigma-Aldrich (Deisenhofer, Germany). Methanol, dimethyl sulfoxide (DMSO), hexane and butylhydroxytoluene (BHT) and culture media (including Mueller-Hinton agar, tryptic soy broth and tryptic soy agar) were purchased from Merck (Germany). *Staphylococcus (S.) aureus* (ATCC 29213) and *Escherichia (E.) coli* O157:H7 (ATCC 35218) were provided by the Iranian Research Organization for Science and Technology (Iran).

2.2. Date seeds

Four Iranian date varieties (Kabkab, Rabbi, Zahedi and Mazafati) were provided by the Iranian Date Palm and Tropical Fruits Research Center, Ahwaz, Iran. Date seeds were removed from the fruits and washed to remove impurities. Pits were dried in oven at 40°C for 24 h and then completely grounded to make fine powders using laboratory scale mortar grinder (Pulverisette, Fritsch, Germany) [21]. These powders were stored at -40°C in deep freezer (Media, WHS-109 FW1, USA) until use [22].

2.3. Preparation of the date palm pit extracts

Extracts of date seeds were prepared based on a previously described method by Al-Farsi and Lee with modifications [23]. Briefly, 1 g of each date pit powder was soaked in 20 ml of hexane. After rinsing, these were extracted at 40°C for 2 h using 60 ml of 80% ethanol and stirring rate of 120 rpm. During the extraction process, conical flasks were covered with aluminum foils to prevent solvent evaporation and light-induced changes. Mixtures were centrifuged at 3913 ×g for 20 min at room temperature. Supernatants were then filtered through Whatman No. 1 filter papers. Filtered extracts were concentrated to dryness at 40°C and reduced pressure using rotary evaporator (HS 2005s Hahnshin Scientific, South Korea). The remaining extracts were stored at -20°C until use.

2.4. Assessment of total phenolic content

Phenolic contents of the date seed extracts were assessed based on the method of Al-Farsi and Alasalvar [24]. Briefly, 200 µl of the seed extracts were mixed with 1.5 ml of Folin-Ciocalteu reagent (10-fold diluted). After incubation of the mixtures at 22°C for 5 min, 1.5 ml of sodium bicarbonate solution (60 g l⁻¹) were added to the mixture. After 90 min incubation at 22°C, absorbance of the mixtures was measured at 725 nm using UV-Vis Absorption 1601 Spectrophotometer (Shimadzu, Japan). Total phenolic contents were assessed using calibration curve (R² = 0.99) of gallic acid as standard (0.001-0.006 mg ml⁻¹). Total phenolic contents of the extracts were expressed as mg of gallic acid equivalent per 100 g of dry weight of the date seeds (mg GAE 100 g⁻¹ dw).

2.5. Assessment of antioxidant activity

Antioxidant activity of the seed extracts was assessed using DPPH free radicals scavenging method based on a method described by Bakhtiyari et al. [25]. Various concentrations of each extract and BHT (as reference compound) were prepared in DMSO. After adding 50 µl of the prepared solutions to microplates, 200 µl of DPPH in methanol (100 µM) were added to microplates. Reaction mixtures were stored at darkness for 30 min at room temperature. Absorbance of the mixtures was read immediately at 517 nm using BioTek XS2 PowerWave Microplate Reader (BioTek Instruments, USA). Control included 50 µl of DMSO and DPPH solutions and blank included pure methanol. The percentage of inhibition for each extract was calculated based on the following equation 1:

$$\text{Inhibition (\%)} = [1 - (A_{\text{extract}} - A_{\text{blank}}) / A_{\text{control}}] \times 100 \quad \text{Eq. 1}$$

Each extract was prepared in three replicates and the mean values of data were used for the calculation of IC₅₀.

The IC₅₀ values of antioxidants were calculated from the calibration curve [26].

2.6. Analysis of phenolics by HPLC-RP

Phenolic compounds were analyzed using HPLC-RP system with UV-photodiode array detector (Smartline 2800, Knauer, Germany). Compounds were separated on a C-18 reverse phase column (Waters Spherisorb S5 ODS2, 250 × 4.6 mm, particle size of 5 µm). The HPLC profiles of phenolic compounds were monitored at 254, 275, 305 and 320 nm. Injection volume of the extract included 20 µl. The mobile phase included methanol (99.5% v v⁻¹) with 0.05% of trifluoroacetic acid (TFA) as Eluent A and H₂O with 0.05% of TFA as Eluent B. The flow rate included 0.5 ml min⁻¹ and the gradient conditions were programmed as 0–10 min, 5% A; 10–20 min, 5–25% A; 20–30 min, 25–35% A; 30–48 min, 35–45% A; 48–50 min, 45–90% A; 50–52 min, 90–95% A; 52–70 min; 95–5% A; and 70–80 min, 5–100% A. Total time for the analysis of samples included 80 min [27]. Method for the quantification of phenolic compounds included external standard procedure. Used standards included gallic, cinnamic, 3,4-dihydroxybenzoic (DHB), vanillic, chlorogenic, caffeic, and 2,5-dihydroxybenzoic acid. Six various concentrations of the standard stock solutions were prepared and injected to HPLC and then peak areas of chromatograms were plotted against concentrations of the stock solutions to prepare calibration curves. These curves were used to calculate concentrations of polyphenols in the samples. For each calibration curve, the best fit line was determined using linear regression analysis. Contents of the identified polyphenols were calculated from the regression equations of the respective curves. The correlation of coefficient (r) for the regression equations included 0.993–0.997.

2.7. Assessment of antibacterial activity

2.7.1. Bacterial strains

Antibacterial activity of the date seed extracts was assessed against *S. aureus* and *E. coli* O157:H7. Bacterial suspensions from overnight culture at 37°C were prepared in tryptic soy broth. Turbidity of the bacterial cultures was adjusted to 0.5 McFarland standard (1.0 × 10⁸ CFU ml⁻¹) [28].

2.7.2. Disc diffusion method

Antibacterial activity of the extracts against *S. aureus* and *E. coli* O157:H7 was assessed using disc diffusion method [28,29]. All extracts were prepared in sterile conditions and dissolved in distilled water to prepare concentrations of 200 mg ml⁻¹. These were sterilized through Millipore membrane filters with pore sizes of 0.45 µm. Then, 10 µl of the prepared extracts were dropped onto sterile paper discs (6 mm in diameter). The solvent control (water) did not show any antimicrobial activity. Commercial discs of ampicillin (10 µg) were used as

positive control. Then, 100 µl of the overnight incubated bacterial cultures (10⁸ CFU ml⁻¹) were spread over the Muller Hinton agar plates. Discs were placed on the inoculated plates and incubated at 37°C for 24 h. Zones of inhibition were measured and expressed in millimeters.

2.7.3 Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

Based on the results from disk diffusion method, MIC of the seed extracts against *S. aureus* was assessed using broth microdilution technique described by Hammer et al. and Mahmoudzadeh et al. [30,31]. Stock solutions of the extracts (200 mg ml⁻¹) were prepared in 5% of DMSO and sterilized through 0.45 µm filter papers. From these stock solutions, double serial dilutions in tryptic soy broth (100–0.2 mg ml⁻¹) were prepared using microtiter trays with 96 U-bottom wells. Bacteria cultures at a final concentration of 10⁸ CFU ml⁻¹ were inoculated into each well containing 400 µl of the dilutions. Wells containing bacteria in media and without extracts were considered as positive and those containing 5% of DMSO as negative controls. Bacterial growth was assessed by measuring turbidity at 600 nm using Bioscreen C FB-1100 (MBR, Helsinki, Finland) after incubation at 37°C for 24 h. For MBC, 10 µl of broth from each well with no visible bacterial growth were streaked on Mueller-Hinton agar plates. After incubation at 37°C for 24 h, the lowest concentration of the extracts that completely inhibited the bacterial growth was considered as MBC.

2.8. Statistical analysis

All analyses were carried out in three replicates and expressed as means ±SD (standard deviation). One way of analysis of variance was used to compare the groups and significance of differences was reported using Duncans post hoc test at P ≤ 0.05. All the analyses were carried out using SPSS Software v.10.1 (SPSS, Chicago, IL).

3. Results and discussion

3.1. Assessment of total phenolic content

Phenolic contents of the date seed extracts were assessed using Folin-Ciocalteu method. This method has been chosen because of its low-cost, simplicity, repeatability and rapidity [32]. Calculated phenolic contents for the date pit extracts ranged 1483–3377 mg GAE 100 g⁻¹ dw (Fig. 1). The highest phenolic content was observed in Kabkab variety seeds (3377 ±0.45 mg GAE 100 g⁻¹ dw) followed by Zahedi (3310 ±0.27 mg GAE 100 g⁻¹ dw), Rabbi (2423 ±0.50 mg GAE 100 g⁻¹ dw) and Mazafati (1483 ±0.60 mg GAE 100 g⁻¹ dw). Based on Marshall et al. findings [33], total phenolic contents of three date seed varieties supplied in the UK market (namely Deglet Nour, Khouat Allig and Zahedi) ranged 2058–2983 mg GAE 100 g⁻¹ w. In another study, Al Farsi et al. [13] studied the phenolic contents of syrups and byproducts (press cakes and seeds) of three Omani date

varieties. Results showed that the total phenolic content of seeds (3102-4430 mg GAE 100 g⁻¹ w) was higher than those found in date syrups and byproducts (press cakes). However, the phenolic content reported in the current study was higher than those reported in Mansouri et al. [21] study (661 and 572 mg GAE 100 g⁻¹ dw for Deglet Noor and Medjool varieties, respectively) and Juhaimi and Ghafoor [34] (1.98 and 4.65 mg GAE 100 g⁻¹ dw for Barhi and Soughi varieties, respectively). Variations in results are mainly linked to date varieties, extraction methods, physiological conditions of fruits (e.g. stage of ripening) and growing conditions (e.g. growing seasons, soil fertilization and cultivation place) [13,35]. It can be suggested that date seeds are good sources of phenolic compounds including phenolic acids and their derivatives.




Figure 1. Total phenolic contents of four Iranian date palm seed extracts. Values are expressed as the means of three replicates and error bars represent standard deviation (SD) (n = 3). Means with a different superscript letter in each column are significantly different at P≤0.05

3.2. Phenolic profiles of the extracts

Seven phenols of chlorogenic acid, caffeic acid, vanillic acid, gallic acid, cinnamic acid, 3,5-DHB and 2,5-DHB were identified and quantified in four varieties of date seeds as listed in Table 1. Results of HPLC analysis demonstrated all phenolic compounds in the extracts. Fig. 2. Shows selected chromatograms of cinnamic acids; the most abundant phenolic compound found in date seed extracts. Concentrations of the phenolic compounds in the extracts varied from one variety to another one. Cinnamic acid was the most abundant phenolic compound found in date seed extracts with the total concentration of 1.54-3.53 mg g⁻¹ dw; respectively followed by chlorogenic acid (0.17-0.28 mg g⁻¹ dw), caffeic acid (0.17-0.26 mg g⁻¹ dw), 3,5-DHB (0.15-0.27 mg g⁻¹ dw), vanillic acid (0.11-0.2 mg g⁻¹ dw) and gallic acid (0.11-0.18 mg g⁻¹ dw). The least phenol quantity belonged to 2,5-DHB with concentrations ranged 0.1-0.17 mg g⁻¹ dw. The highest content of phenols belonged to Zahedi seed extract; followed by Kabkab seed extract. In contrast, Mazafati and Rabbi seed extracts included the lowest quantities of phenols with total contents of 2.4 and 2.8 mg g⁻¹ dw, respectively. Mansouri et al. [21] reported a similar phenolic profile from seven Algerian date pits. They found that cinnamic acid and its derivatives (di- hydrocinnamic, coumaric and sinapic acids) were the major simple phenols presented in all varieties. Al Juhaimi and Ozcan [36] in a study reported six phenolic compounds including caffeic acid, protocatechuic acid, gallic acid, rutin, catechin and syringic acid in seeds of 12 date varieties from five countries. Similarly, Al-Farsi and Lee [23] found nine phenolic compounds in seeds of Omani dates. In general, p-hydroxybenzoic, protocatechuic and m-coumaric acids were three major compounds, respectively. Contrary to the findings of previous studies, detected hydroxybenzoic compounds in the current study included the least abundant phenolic acids (Table 1).

Table 1. Phenolic profiles of the four Iranian date seed extracts


	Concentration(mg g ⁻¹ dw)							Sum of polyphenols
	Gallic acid	Vanillic acid	3,4 DHB ^a	2,5 DHB ^b	Cinnamic acid	Caffeic acid	Chlorogenic acid	
Rabbi	0.12 ±0.08 ^c	0.13 ±0.09 ^c	0.15±0.01 ^b	0.10 ±0.01 ^c	1.54 ±0.09 ^b	0.17 ±0.05 ^b	0.20 ±0.01 ^c	2.41 ^c
Zahedi	0.18 ±0.13 ^a	0.20 ±0.01 ^a	0.27 ±0.02 ^a	0.17 ±0.09 ^a	3.36 ±0.04 ^d	0.26 ±0.06 ^d	0.28 ±0.01 ^a	4.72 ^a
Kabkab	0.16 ±0.01 ^b	0.15 ±0.02 ^b	0.26 ±0.07 ^a	0.14 ±0.05 ^b	3.53 ±0.05 ^c	0.19 ±0.08 ^c	0.17 ±0.09 ^b	4.6 ^b
Mazafati	0.11 ±0.01 ^d	0.11 ±0.03 ^d	0.15 ±0.01 ^b	0.10 ±0.04 ^c	1.90 ±0.03	0.22 ±0.05 ^a	0.21 ±0.01 ^c	2.8 ^d

Results are expressed as mean ±SD (standard deviation) (n = 3). Means with a different lowercase letter in each row are significantly different at P≤0.05


^a3,4-dihydroxybenzoic acid

^b2,5-dihydroxybenzoic acid

A



B






Figure 2. Selected chromatograms from RP-HPLC analysis of four date palm seed extracts. A) Kabkab; B) Zahedi; C) rabbi and D) Mazafati

3.3. Assessment of antioxidant activity

Antioxidant activity of the date seed extracts and BHT are presented in Fig. 3. The IC_{50} values of the seed extracts varied 16.55–21.58 $\mu\text{g ml}^{-1}$. Kabkab seed extracts included the highest antioxidant activity against DPPH radicals; followed by Rabbi (20.4 $\mu\text{g ml}^{-1}$) and Zahedi (20.8 $\mu\text{g ml}^{-1}$) seed extracts. Mazafati variety with a high IC_{50} of 21.58 $\mu\text{g ml}^{-1}$ included the least antioxidant activity. However, the radical scavenging effect of all extracts was less than that of BHT, but the IC_{50} value of Kabkab variety was close to that of BHT as standard reference. It is noteworthy that pure BHT was more active than crude BHT [37]. To assess the contribution of total phenolic contents to antioxidant capacity of the date seeds, Pearson's correlation coefficient was calculated. The analysis of correlation ($P < 0.05$) showed a significant negative correlation between the IC_{50} and the total phenolic content of the date seed extracts ($r = -0.75$). Based on previously published studies, correlations have been reported between the antioxidant activity and the total phenolic content of dates [35,38].




Figure 3. DPPH radical scavenging activities (IC_{50} $\mu\text{g ml}^{-1}$) of four Iranian date palm seed extracts. Values are expressed as the means of three replicates and error bars represent standard deviation (SD) ($n = 3$). Means with a different superscript letter in each column are significantly different at $P \leq 0.05$.

Studies have reported that seeds from various fruits including mango, basil-seed, wild pear, pomegranate and red grape include antioxidant activities [39–43]. Antioxidant activity of the date seed extracts is mainly due to the presence of phenolic compounds. Results of several studies have demonstrated that phenolic compounds such as cinnamic acid and its derivatives (caffeic acid, p-coumaric acid, ferulic acid, chlorogenic acid and sinapic acid) offer a higher antioxidative capacity, compared to other polyphenols such as hydroxybenzoic acid derivatives (gallic acid, vanillic acid, dihydroxybenzoic acids and protocatechuic acid). Considering that date seeds include considerable quantities of these compounds (especially cinnamic acid derivatives), it can be concluded that date seeds are rich sources of antioxidant compounds [44–46].

3.4. Assessment of antibacterial activity

In the current study, antibacterial activity of the ethanolic extracts of date seeds was assessed against *E. coli* O157:H7 as Gram-negative and *S. aureus* as Gram-positive bacteria (Table 2). Findings from the disc diffusion method revealed that all evaluated seed extracts included varying antibacterial effects against *S. aureus* but not against *E. coli*. Differences in susceptibilities of the two bacteria against date seed extracts could be attributed to the differences in structure of the bacterial cell walls. Cytoplasm of the Gram-positive bacteria is surrounded by a thick peptidoglycan layer, while in Gram-negative bacteria an additional outer membrane mainly containing lipopolysaccharides is present [47,48]. Antimicrobial strength of the phenolic compounds against Gram-positive bacteria is associated with their ability to interact directly with the peptidoglycan layer and thereby to decrease cell integrity and to increase bacterial sensitivity to osmotic pressure and ionic strength. However, the lipopolysaccharide layer in outer membrane of Gram-negative bacteria, which functions as a tight permeability barrier, hinders polyphenols to have direct connections with the peptidoglycan layer [49].

Regarding antibacterial effects of the highlighted extracts against *S. aureus*, the maximum effect was observed for Kabkab variety (12.2 mm), followed by Zahedi (11.5 mm), Rabbi (10 mm) and Mazafati (0.9 mm) varieties. Differences in antibacterial activities of the extracts could be attributed to the differences in their qualitative compositions. As previously shown in analysis of phenolic compounds, the highlighted seed extracts included the highest quantity of cinnamic acid and its derivatives (including chlorogenic acids). These compounds in comparison with other phenolics such as hydroxyl benzoic acids include notable antibacterial activity, especially against Gram-positive bacteria such as *S. aureus* [48]. The antibacterial potency of these extracts against *S. aureus* was subsequently assessed using MIC and MBC methods (Table 3). Kabkab and Zahedi date extracts included the highest inhibition activity or lowest MIC against *S. aureus* (1.56 mg ml^{-1}) followed by Rabbi and Mazafati dates (3.125 mg ml^{-1}). Furthermore, MBC method showed that all investigated extracts included antibacterial activity. Despite the fact that a similar inhibitory effect against *S. aureus* was reported for Kabkab and Zahedi dates, the bactericidal effect of Kabkab date was more than that of Zahedi date (the lowest MBC of 3.125 mg ml^{-1}). In contrast to Kabkab date, Mazafati and Rabbi date seed extracts demonstrated the weakest antibacterial activity (MBC of 12.50 mg ml^{-1}).

Table 2. Antibacterial activities of the four date seed extracts using disc diffusion method

	Zahedi	Kabkab	Rabbi	Mazafati	Ampicillin
	Inhibition zone (mm) ^a				
<i>Staphylococcus aureus</i>	11.5 ±0.06 ^b	12.2 ±0.05 ^b	10.0 ±0.2 ^a	0.9 ±0.05 ^a	18 ±0.25 ^c
<i>Escherichia coli</i>	ND	ND	ND	ND	21 ±0.5

Results are expressed as mean ±SD (standard deviation) ($n = 3$); ND, not detected; means with a different lowercase letter in each row are significantly different at $P \leq 0.05$

^aInhibition zone diameter (mm)

Table 3. Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of the four date seed extracts against *Staphylococcus aureus* (mg m^{-1})

	Zahedi		Kabkab		Rabbi		Mazafati	
	MIC (mg m^{-1})	MBC (mg m^{-1})	MIC (mg m^{-1})	MBC (mg m^{-1})	MIC (mg m^{-1})	MBC (mg m^{-1})	MIC (mg m^{-1})	MBC (mg m^{-1})
<i>Staphylococcus aureus</i>	1.56 ^b	6.25 ^C	1.56 ^b	3.125 ^B	3.125 ^a	12.5 ^A	3.125 ^a	12.5 ^A

Results are expressed as mean ±SD (standard deviation) ($n = 3$)

Different lowercase and capital letters respectively indicate significant differences between MIC and MBC of the four date seed extracts ($P \leq 0.05$)

MIC=Minimum inhibitory concentration

MBC= minimum bactericidal concentration

According to Qadoos et al. [50], date fruit and leave extracts show inhibitory effects against Gram-positive bacteria (*S. aureus* and *B. subtilis*); however, no effects are seen for Gram-negative bacteria (*E. coli* and *Pseudomonas aeruginosa*). Samad et al. [51] studied effects of methanolic extracts of various date fruits against Gram-positive and Gram-negative bacteria (*Serratia marcescens* and *E. coli*) and reported that the significant differences against studied bacteria were associated to various concentrations of methanolic extracts (100-500 mg ml^{-1}) from date varieties. Of these varieties, Ajwa dates with the highest quantities of phenolic compounds showed antimicrobial activities against all four bacteria at higher concentrations; however, no inhibitory activities were observed against Gram-negative bacteria in Mabroom and Mariami study. Based on the results from the current study, increased phenolic contents can result in increased antimicrobial activity of the extracts with a correlation factor of 0.99. Several studies have shown that antimicrobial activity of the extracts is directly linked to their polyphenolic compounds [52-55]. The mechanism of action of these compounds in inhibiting the bacterial growth can be summarized as interaction with the bacterial cell walls [54,56], precipitation of proteins and disruption of enzyme activities [52,53,55,57,58].

4. Conclusion

This study demonstrated that all the assessed extracts included significant phenolic contents, mainly cinnamic acid and its derivatives, as well as antioxidant activities. The best results were observed for Kabkab date seed extracts. Moreover, the study revealed that all palm date seeds included antibacterial activities against *S. aureus* but not against *E. coli*. In general, palm date seeds (as byproducts of the processing industries) are rich sources of phenolic compounds, which can be used as complementary

biotechnological substrates or as nutraceuticals for the production of novel, healthy innovative foods. However, further human clinical studies seem necessary.

5. Acknowledgements

This study was supported financially by the National Nutrition and Food Technology Research Institute (NNFTRI) of Iran.

6. Conflict of Interest

The authors declare no conflict of interest.

References

- Mohamed RM, Fageer AS, Eltayeb MM, Mohamed Ahmed IA. Chemical composition, antioxidant capacity, and mineral extractability of Sudanese date palm (*Phoenix dactylifera L.*) fruits. *Food Sci Nutr.* 2014; 2(5): 478-489
doi: 10.1002/fsn3.123
- Johnson DV, Al-Khayri JM, Jain SM. Introduction: Date Production Status and Prospects in Africa and the Americas. *Date Palm Genetic Resources and Utilization*: Springer; 2015. pp. 3-18.
doi: 10.1007/978-94-017-9694-1_1
- Zhang CR, Aldosari SA, Vidyasagar PS, Shukla P, Nair MG. Health-benefits of date fruits produced in Saudi Arabia based on in vitro antioxidant, anti-inflammatory and human tumor cell proliferation inhibitory assays. *J Saud Soc Agric Sci.* 2017; 16(3): 287-293.
doi: 10.1016/j.jssas.2015.09.004
- Faostat F. Agriculture Organization of the United Nations statistics division (2014). Production available in: <http://faostat3.fao.org/browse/Q/QC/S> [Review date: April 2015]. 2016
- Besbes S, Blecker C, Deroanne C, Lognay G, Drira N, Attia H. Quality characteristics and oxidative stability of date seed oil during storage. *Food Sci Technol Int.* 2004; 10(5): 333-338.
doi: 10.1177/1082013204047777

6. Besbes S, Drira L, Blecker C, Deroanne C, Attia H. Adding value to hard date (*Phoenix dactylifera* L.): Compositional, functional and sensory characteristics of date jam. *Food Chem.* 2009; 112(2): 406-411. doi: 10.1016/j.foodchem.2008.05.093
7. Khiyami M, Aboseide B, Pometto A. Influence of complex nutrient sources: Dates syrup and dates pits on *Lactococcus lactis* growth and nisin production. *J Biotechnol.* 2008; 136: S736. doi: 10.1016/j.jbiotec.2008.07.1754
8. Abou Zeid AA, Baeshin NA, Baghlaf AO. Utilization of date products in production of oxytetracycline by *Streptomyces rimosus*. *Zbl Mikrobiol.* 1993; 148(5): 333-341. doi: 10.1016/S0232-4393(11)80122-8
9. Al-Ogaidi H, Khalifa S, Hadi H, Al-Nakash S. Production of protein from date stone by *Aspergillus oryzae*. *J Agric Water Resour Res.* 1985; 4(3): 197-206
10. Abou Zeid AZA, Baghlaf AO, Khan JA, Makhashin SS. Utilization of date seeds and cheese whey in production of citric acid by *Candida lipolytica*. *Agr Wastes.* 1983;8(3): 131-142. doi: 10.1016/0141-4607(83)90112-9
11. Hossain MZ, Waly MI, Singh V, Sequeira V, Rahman MS. Chemical composition of date-pits and its potential for developing value-added product-a review. *Pol J Food Nutr Sci.* 2014; 64(4): 215-226. doi: 10.2478/pjfn-2013-0018
12. Hamada J, Hashim I, Sharif F. Preliminary analysis and potential uses of date pits in foods. *Food Chem.* 2002; 76(2): 135-137. doi: 10.1016/S0308-8146(01)00253-9
13. Al-Farsi M, Alasalvar C, Al-Abid M, Al-Shoaily K, Al-Amry M, Al-Rawahy F. Compositional and functional characteristics of dates, syrups, and their by-products. *Food Chem.* 2007; 104(3): 943-947. doi: 10.1016/j.foodchem.2006.12.051
14. Sahari M, Barzegar M, Radfar R. Effect of varieties on the composition of dates (*Phoenix dactylifera* L.)- note. *Food Sci Technol Int.* 2007; 13(4): 269-275. doi: 10.1177/1082013207082244
15. Bailey G, Williams D. Potential mechanisms for food-related carcinogens and anticarcinogens. *Food Technol.* 1993; 47(2): 105-118
16. Mohamed Lemine FM, Mohamed Ahmed MVO, Ben Mohamed Maoulainine L, Bouna ZeAO, Samb A, Boukhary AOMS. Antioxidant activity of various mauritanian date palm (*Phoenix dactylifera* L.) fruits at two edible ripening stages. *Food Sci Nutr.* 2014; 2(6): 700-705. doi: 10.1002/fsn3.167
17. Liverio L, Puglisi P, Morazzoni P, Bombardelli E. Antimutagenic activity of procyanidins from *vitis vinifera*. *J Med Plants Stud.* 1994; 65: 203-209
18. Takechi M, Tanaka Y, Takehara M, Nonaka GI, Nishioka I. Structure and antiherpetic activity among the tannins. *Phytochemistry.* 1985; 24(10): 2245-2250. doi: 10.1016/S0031-9422(00)83018-6
19. Farag MA, Mohsen M, Heinke R, Wessjohann LA. Metabolomic fingerprints of 21 date palm fruit varieties from Egypt using UPLC/PDA/ESI-qTOF-MS and GC-MS analyzed by chemometrics. *Food Res Int.* 2014; 64: 218-226. doi: 10.1016/j.foodres.2014.06.021
20. Basanta MF, Rizzo SA, Szerman N, Vaudagna SR, Descalzo AM, Gerschenson LN, Perez cd, Rojas AM. Plum (*Prunus salicina*) peel and pulp microparticles as natural antioxidant additives in breast chicken patties. *Food Res Int.* 2018; 106: 1086-1094. doi: 10.1016/j.foodres.2017.12.011
21. Mansouri A, Embarek G, Kokkalou E, Kefalas P. Phenolic profile and antioxidant activity of the algerian ripe date palm fruit (*Phoenix dactylifera*). *Food Chem.* 2005; 89(3): 411-420. doi: 10.1016/j.foodchem.2004.02.051
22. Singh V, Guizani N, Essa M, Hakkim F, Rahman M. Comparative analysis of total phenolics, flavonoid content and antioxidant profile of different date varieties (*Phoenix dactylifera* L.) from Sultanate of Oman. *Int Food Res J.* 2012; 19(3): 1063
23. Al-Farsi MA, Lee CY. Optimization of phenolics and dietary fibre extraction from date seeds. *Food Chem.* 2008; 108(3): 977-985. doi: 10.1016/j.foodchem.2007.12.009
24. Al-Farsi M, Alasalvar C, Morris A, Baron M, Shahidi F. Comparison of antioxidant activity, anthocyanins, carotenoids, and phenolics of three native fresh and sun-dried date (*Phoenix dactylifera* L.) varieties grown in Oman. *J Agric Food Chem.* 2005; 53(19): 7592-7599. doi: 10.1021/jf050579q
25. Bakhtiyari E, Ahmadian Attari MM, Salehi P, Khallaghi B, Dargahi L, Mohamed Z, Ahmadiani A, Kamalinejad M. Non-polyphenolic compounds of a specific kind of dried grape (Maviz) inhibit memory impairments induced by beta-amyloid peptide. *Nutr Neurosci.* 2017; 20(8): 469-477. doi: 10.1080/1028415X.2016.1183986
26. Sirisena S, Ajlouni S, Ng K. Simulated gastrointestinal digestion and In vitro colonic fermentation of date (*Phoenix dactylifera* L.) seed polyphenols. *Int J Food Sci Technol.* 2018; 53(2): 412-422. doi:10.1111/ijfs.13599
27. Moradi Afrapoli F, Asghari B, Saeidnia S, Ajani Y, Mirjani M, Malmir M, Bazaz RD, Hadjiakhoondi A, Salehi P, Hamburger M, Yassa N. In vitro α -glucosidase inhibitory activity of phenolic constituents from aerial parts of *Polygonum hyrcanicum*. *DARU J Pharm Sci.* 2012; 20(1): 20-37. doi: 37.10.1186/2008-2231-20-37
28. Abdollahzadeh E, Rezaei M, Hosseini H. Antibacterial activity of plant essential oils and extracts: The role of thyme essential oil, nisin, and their combination to control *Listeria monocytogenes* inoculated in minced fish meat. *Food Control.* 2014; 35(1): 177-183. doi: 10.1016/j.foodcont.2013.07.004
29. Bauer A, Kirby W, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol.* 1966; 45(4): 493. doi: 10.1093/ajcp/45.4_ts.493
30. Hammer KA, Carson CF, Dunstan JA, Hale J, Lehmann H, Robinson CJ, Prescott SL, Riley TV. Antimicrobial and anti-inflammatory activity of five *Taxandria fragrans* oils In vitro. *Microbiol Immunol.* 2008; 52(11): 522-530. doi: 10.1111/j.1348-0421.2008.00070.x
31. Mahmoudzadeh M, Hosseini H, Shahraz F, Akhondzadeh Basti A, Khaneghah AM, Azizkhani M, Sant'ana ADS, Haghshenas M, Mahmoudzadeh L. Essential oil composition

- and antioxidant capacity of *carum copticum* and its antibacterial effect on *Staphylococcus aureus*, *Enterococcus faecalis* and *Escherichia coli* O157: H7. *J Food Process Preserv.* 2017; 41(3): e12938.
doi: 10.1111/jfpp.12938
32. Bouhlali E, Bammou M, Sellam K, Benlyasa M, Alem C, Filali Zegzouti Y. Evaluation of antioxidant, antihemolytic and antibacterial potential of six Moroccan date fruit (*Phoenix dactylifera*L.) varieties. *J King Saud Univ Sci.* 2016; 28: 136-142.
doi: 10.1016/j.jksus.2016.01.002
 33. Marshall R, Mistrello J, Sirisena S, Ghavami A, Krishnamoorthy S. Determination of the antioxidant capacity, total phenolic and avonoid contents of seeds from three commercial varieties of culinary dates. *Int J Food Stud.* 2014; 3(1): 34-44.
doi: 10.7455/ijfs/3.1.2014.a3
 34. Juhaimi FA, Ghafoor K, Ozcan MM. Physical and chemical properties, antioxidant activity, total phenol and mineral profile of seeds of seven different date fruit (*Phoenix dactylifera* L.) varieties. *Int J Food Sci Nutr.* 2012; 63(1): 84-89.
doi: 10.3109/09637486.2011.598851
 35. Ardekani MRS, Khanavi M, Hajimahmoodi M, Jahangiri M, Hadjiakhoondi A. Comparison of antioxidant activity and total phenol contents of some date seed varieties from Iran. *Iran J Pharm Res.* 2010; 9(2): 141-146
 36. Al Juhaimi F, Ozcan MM, Adiamo OQ, Alsawmahi ON, Ghafoor K, Babiker EE. Effect of date varieties on physico-chemical properties, fatty acid composition, tocopherol contents, and phenolic compounds of some date seed and oils. *J Food Process Preserv.* 2018; 42(4): e13584.
doi: 10.1111/jfpp.13584
 37. Kokabi M, Yousefzadi M, Fegghi MA, Keshavarz M. Antioxidant activity of extracts of selected algae from the persian gulf, Iran. *J Persian Gulf.* 2013; 4(12): 45-50
 38. Biglari F, AlKarkhi AF, Easa AM. Antioxidant activity and phenolic content of various date palm (*Phoenix dactylifera*) fruits from Iran. *Food Chem.* 2008; 107(4): 1636-1641.
doi: 10.1016/j.foodchem.2007.10.033
 39. Soong YY, Barlow PJ. Antioxidant activity and phenolic content of selected fruit seeds. *Food Chem.* 2004; 88(3): 411-417.
doi: 10.1016/j.foodchem.2004.02.003
 40. Hashemi SMB, Mousavi Khaneghah A. Characterization of novel basil-seed gum active edible films and coatings containing oregano essential oil. *Prog Org Coat.* 2017; 110: 35-41.
doi: 10.1016/j.porgcoat.2017.04.041
 41. Hashemi SMB, Khaneghah AM, Barba FJ, Lorenzo JM, Rahman MS, Amarowicz R, Yousefabad SA, Dabiri Movahed M. Characteristics of wild pear (*Pyrus glabra* Boiss) seed oil and Its oil- in- water emulsions: A novel source of edible oil. *Eur J Lipid Sci Technol.* 2018; 120(2): 1700284.
doi: 10.1002/ejlt.201700284
 42. Derakhshan Z, Ferrante M, Tadi M, Ansari F, Heydari A, Hosseini MS, Contid GO, Khalili Sadrabad E. Antioxidant activity and total phenolic content of ethanolic extract of pomegranate peels, juice and seeds. *Food Chem Toxicol.* 2018; 114: 108-111.
doi: 10.1016/j.fct.2018.02.023
 43. Hashemi SMB, Mousavi Khaneghah A, Nikmaram N, Raeisi S, Rahman MS, Avallone S. Heating, microwave and UV irradiation effects on oxidative stability of Sardasht red grape (*Vitis vinifera* cultiv. Sardasht) seed oil. *Int J Food Sci Technol.* 2017; 52(6): 1341-1347.
doi: 10.1111/ijfs.13421
 44. Connor AM, Luby JJ, Hancock JF, Berkheimer S, Hanson EJ. Changes in fruit antioxidant activity among blueberry cultivars during cold-temperature storage. *J Agric Food Chem.* 2002; 50(4): 893-898.
doi: 10.1021/jf011212y
 45. Guo C, Yang J, Wei J, Li Y, Xu J, Jiang Y. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr Res.* 2003; 23(12): 1719-1726.
doi: 10.1016/j.nutres.2003.08.005
 46. Cuvelier ME, Richard H, Berset C. Comparison of the antioxidative activity of some acid-phenols: Structure-activity relationship. *Biosci Biotechnol Biochem.* 1992; 56(2): 324-325.
doi: 10.1271/bbb.56.324
 47. Alzoreky N, Nakahara K. Antibacterial activity of extracts from some edible plants commonly consumed in Asia. *Int J Food Microbiol.* 2003; 80(3): 223-230.
doi: 10.1016/S0168-1605(02)00169-1
 48. Fattouch S, Caboni P, Coroneo V, Tuberoso CIG, Angioni A, Dessi S, Marzouki N, Cabras P. Antimicrobial activity of tunisian quince (*cydonia oblonga* miller) pulp and peel polyphenolic extracts. *J Agric Food Chem.* 2007; 55(3): 963-969.
doi: 10.1021/jf062614e
 49. Papuc C, Goran GV, Predescu CN, Nicorescu V, Stefan G. Plant polyphenols as antioxidant and antibacterial agents for shelf-life extension of meat and meat products: Classification, structures, sources, and action mechanisms. *Compr Rev Food Sci Food Saf.* 2017; 16(6): 1243-1268.
doi:10.1111/1541-4337.12298
 50. Qadoos HA, Dhafari HS, Al Marzooqi DA, Kumarappan A, Nazir A. Phenolic content and antimicrobial activities of date palm (*Phoenix dactylifera* L.) fruits and leaves. *Food Biol.* 2017; 6: 11-15.
doi: 10.19071/fbiol.2017.v6.3154
 51. Samad MA, Hashim SH, Simarani K, Yaacob JS. Antibacterial properties and effects of fruit chilling and extract storage on antioxidant activity, total phenolic and anthocyanin content of four date palm (*Phoenix dactylifera*) cultivars. *Molecules.* 2016; 21(4): 419.
doi: 10.3390/molecules21040419.
 52. Asgharzadeh A, Aubourg SP, Hosseini H, Shabanpour B. Chemical changes in silver carp (*Hypophthalmichthys molitrix*) minced muscle during frozen storage: Effect of a previous washing process. *Grasas Aceites.* 2010; 61: 95-101
doi: 10.3989/gya.087109
 53. Bakhtiary F, Sayevand HR, Khaneghah AM, Haslberger AG, Hosseini H. Antibacterial efficacy of essential oils and sodium nitrite in vacuum processed beef fillet. *Appl food Biotechnol.* 2018; 5(1): 1-10.
doi: 10.22037/afb.v5i1.17118
 54. Barbary O, El Sohaimy S, El Saadani M, Zeitoun A. Antioxidant, antimicrobial and anti-HCV activities of lignan extracted from flaxseed. *Res J Agric Biol Sci.* 2010; 6(3): 247-256

55. Papuc C, Goran GV, Predescu CN, Nicorescu V, Stefan G. Plant polyphenols as antioxidant and antibacterial agents for shelf- life extension of meat and meat products: Classification, structures, sources, and action mechanisms. *Compr Rev Food Sci Food Saf.* 2017; 16(6): 1243-1268. doi: 10.1111/1541-4337.12298
56. Sellam K, Ramchoun M, Alem C, El Rhaffari L. Biological investigations of antioxidant-antimicrobial properties and chemical composition of essential oil from *Lavandula multifida*. *Oxid Antioxid Med Sci.* 2013; 2(3): 211-216. doi: 10.5455/oams.290513.or.044
57. Eftekhari A, Ahmadian E, Panahi Azar V, Hosseini H, Tabibiazar M, Maleki Dizaj S. Hepatoprotective and free radical scavenging actions of quercetin nanoparticles on aflatoxin B1-induced liver damage: In vitro/In vivo studies. *Artif Cells Nanomed Biotechnol.* 2018; 46(2): 411-420. doi: 10.1080/21691401.2017.1315427
58. Naseri M, Rezaei M, Moieni S, Hosseini H, Eskandari S. Effect of different precooking methods on chemical composition and lipid damage of silver carp (*Hypophthalmichthys molitrix*) muscle. *Int J Food Sci Technol.* 2010; 45(10): 1973-1979. doi:10.1111/j.1365-2621.2010.02349.x

بررسی محتویات فنولی، فعالیت ضداکسایشی و ضد باکتریایی عصاره های هسته چهار رقم خرماي ایرانی (*Phoenix dactylifera L.*)

رامین رادفر^۱، مهدی فرهودی^۱، اسماعیل قاسمی^۲، امین موسوی خانقاه^۳، فرزانه شهرآز^۴، هدایت حسینی^{۱*}

۱- انستیتو تحقیقات تغذیه ای و صنایع غذایی کشور، دانشکده علوم تغذیه و صنایع غذایی، دانشگاه علوم پزشکی شهید بهشتی، تهران، ایران.

۲- گروه پلاستیک، پژوهشگاه پلیمر و پتروشیمی، تهران-ایران.

۳- گروه علوم غذایی، دانشکده مهندسی غذایی، دانشگاه کمپیناس، کمپیناس، اس پی، برزیل.

۴- گروه ریاضیات عمومی و کاربردی، دانشگاه ایالتی صنعت و نفت آذربایجان، باکو، آذربایجان.

چکیده

سابقه و هدف: هر ساله، در صنایع فرآوری و بسته بندی خرما مقادیر زیادی هسته خرما تولید می شوند که یا دور ریخته می شوند یا به عنوان مواد کم ارزش به مصرف خوراک دام و تولید کمیوست می رسند. در حالی که، اینها منابع زیستی^۱ می باشند، که امکان تولید مواد با ارزش افزوده بالا در صنایع غذایی از آنها وجود دارد. هدف اصلی تحقیق حاضر، تعیین پروفایل و محتوای فنولی، فعالیت ضد باکتریایی و ضد اکسایشی عصاره های هسته چهار رقم خرماي ایرانی مضافتی، ربی، کبکاب و زاهدی بوده است.

مواد و روش ها: محتوی فنولی کل، پروفایل ترکیبات فنولی، فعالیت ضداکسایشی و فعالیت ضد باکتریایی عصاره هسته چهار رقم خرماي ایرانی، به ترتیب با استفاده از روش های فولین- سیوکالتو، کروماتوگرافی مایع با کارایی بالا با فاز معکوس، گیرنده ۲، ۲- دی فنیل -۱- پیکریل هیدرازیل، و روش های انتشار دیسک در آگار، تعیین گردیدند.

یافته ها و نتیجه گیری: محتوای فنولی کل بین ۱۴۸۰ تا ۳۳۸۰ میلی گرم بر ۱۰۰ گرم وزن خشک بر حسب اسید گالیک و شامل اسیدهای سینامیک، کلروژنیک، کافئیک و ۳ و ۵- هیدروکسی بنزوئیک به ترتیب عمده ترین ترکیبات فنولی شناسایی شده بودند. در میان ارقام مورد آزمایش، عصاره های هسته ارقام کبکاب و مضافتی با داشتن IC₅₀ به ترتیب معادل ۱۶/۵۶ و ۲۲/۶ کمترین و بیشترین فعالیت گیرندگی رادیکال را به خود اختصاص دادند. نهایتاً یافته های حاصل از آزمون انتشار دیسک نشان داد که کلیه عصاره های هسته خرما روی باکتری *استافیلوکوکوس اورئوس* اثر بازدارندگی داشتند، اما روی باکتری *اشرشیا کلی* تأثیری نداشتند. حداقل غلظت مهاري^۲ و حداقل غلظت کشندگی^۳ عصاره هسته های خرما برای باکتری *استافیلوکوکوس اورئوس* به ترتیب بین ۱/۵۶ تا ۳/۱۲۵ و ۳/۱۲۵ تا ۱۲/۵ میلی گرم بر میلی لیتر تعیین شد. بر اساس یافته های این تحقیق هسته خرماهای ایرانی منبع خوبی برای ترکیبات فنولی قابل استخراج با فعالیت های قایل توجه ضداکسایشی می باشند و از این رو می توانند به عنوان افزودنی های طبیعی در فرمولاسیون فرآورده های گوناگون مانند غذاهای فراسودمند و مکمل های رژیمی مورد استفاده قرار گیرند. علاوه بر این، این هسته ها می توانند از طریق فرایندهای زیست فناوری به فرآورده هایی با ارزش افزوده تبدیل شوند.

تعارض منافع: نویسندگان اعلام می کنند که هیچ نوع تعارض منافی مرتبط با انتشار این مقاله ندارند.

تاریخچه مقاله

دریافت ۱۵ نوامبر ۲۰۱۸

داوری ۲۴ دسامبر ۲۰۱۸

پذیرش ۷ ژوئن ۲۰۱۹

واژگان کلیدی

- خواص ضد باکتریایی
- فعالیت ضداکسایشی
- هسته خرما
- پروفایل فنولی

* نویسنده مسئول

هدایت حسینی،

گروه علوم و صنایع غذایی، انستیتو تحقیقات تغذیه ای و صنایع غذایی کشور، دانشکده علوم تغذیه و صنایع غذایی، دانشگاه علوم پزشکی شهید بهشتی، تهران، ایران

تلفن: +۹۸-۲۱۲۲۳۷۶۴۲۶

نمابر: +۹۸-۲۱۲۲۳۶۰۶۶۰

پست الکترونیک:

hedayat@sbmu.ac.ir

¹ Bioresources

² Minimum inhibitory concentration or MIC

³ Minimum bactericidal concentration or MBC