

Hospital Anxiety and Depression Scale (HADS)

Tick the box beside the reply that is closest to how you have been feeling in the past week.
Don't take too long over your replies: your immediate is best.

D	A		D	A	
		I feel tense or 'wound up':			I feel as if I am slowed down:
	3	Most of the time	3		Nearly all the time
	2	A lot of the time	2		Very often
	1	From time to time, occasionally	1		Sometimes
	0	Not at all	0		Not at all
		I still enjoy the things I used to enjoy:			I get a sort of frightened feeling like 'butterflies' in the stomach:
0		Definitely as much		0	Not at all
1		Not quite so much		1	Occasionally
2		Only a little		2	Quite Often
3		Hardly at all		3	Very Often
		I get a sort of frightened feeling as if something awful is about to happen:			I have lost interest in my appearance:
	3	Very definitely and quite badly	3		Definitely
	2	Yes, but not too badly	2		I don't take as much care as I should
	1	A little, but it doesn't worry me	1		I may not take quite as much care
	0	Not at all	0		I take just as much care as ever
		I can laugh and see the funny side of things:			I feel restless as I have to be on the move:
0		As much as I always could		3	Very much indeed
1		Not quite so much now		2	Quite a lot
2		Definitely not so much now		1	Not very much
3		Not at all		0	Not at all
		Worrying thoughts go through my mind:			I look forward with enjoyment to things:
	3	A great deal of the time	0		As much as I ever did
	2	A lot of the time	1		Rather less than I used to
	1	From time to time, but not too often	2		Definitely less than I used to
	0	Only occasionally	3		Hardly at all
		I feel cheerful:			I get sudden feelings of panic:
3		Not at all		3	Very often indeed
2		Not often		2	Quite often
1		Sometimes		1	Not very often
0		Most of the time		0	Not at all
		I can sit at ease and feel relaxed:			I can enjoy a good book or radio or TV program:
	0	Definitely	0		Often
	1	Usually	1		Sometimes
	2	Not Often	2		Not often
	3	Not at all	3		Very seldom

Please check you have answered all the questions

Scoring:

Total score: Depression (D) _____ Anxiety (A) _____

0-7 = Normal

8-10 = Borderline abnormal (borderline case)

11-21 = Abnormal (case)

Name: _____

Date: _____

Pittsburgh Sleep Quality Index (PSQI)

Instructions: The following questions relate to your usual sleep habits during the past month only. Your answers should indicate the most accurate reply for the majority of days and nights in the past month. **Please answer all questions.**

1. During the past month, what time have you usually gone to bed at night? _____
2. During the past month, how long (in minutes) has it usually taken you to fall asleep each night? _____
3. During the past month, what time have you usually gotten up in the morning? _____
4. During the past month, how many hours of actual sleep did you get at night? (This may be different than the number of hours you spent in bed.) _____

5. During the <u>past month</u> , how often have you had trouble sleeping because you...	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week
a. Cannot get to sleep within 30 minutes				
b. Wake up in the middle of the night or early morning				
c. Have to get up to use the bathroom				
d. Cannot breathe comfortably				
e. Cough or snore loudly				
f. Feel too cold				
g. Feel too hot				
h. Have bad dreams				
i. Have pain				
j. Other reason(s), please describe:				
6. During the past month, how often have you taken medicine to help you sleep (prescribed or "over the counter")?				
7. During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity?				
	No problem at all	Only a very slight problem	Somewhat of a problem	A very big problem
8. During the past month, how much of a problem has it been for you to keep up enough enthusiasm to get things done?				
	Very good	Fairly good	Fairly bad	Very bad
9. During the past month, how would you rate your sleep quality overall?				

	No bed partner or room mate	Partner/room mate in other room	Partner in same room but not same bed	Partner in same bed
10. Do you have a bed partner or room mate?				
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week
If you have a room mate or bed partner, ask him/her how often in the past month you have had:				
a. Loud snoring				
b. Long pauses between breaths while asleep				
c. Legs twitching or jerking while you sleep				
d. Episodes of disorientation or confusion during sleep				
e. Other restlessness while you sleep, please describe:				

Scoring the PSQI

The order of the PSQI items has been modified from the original order in order to fit the first 9 items (which are the only items that contribute to the total score) on a single page. Item 10, which is the second page of the scale, does not contribute to the PSQI score.

In scoring the PSQI, seven component scores are derived, each scored 0 (no difficulty) to 3 (severe difficulty). The component scores are summed to produce a global score (range 0 to 21). Higher scores indicate worse sleep quality.

Component 1: Subjective sleep quality—question 9

<u>Response to Q9</u>	<u>Component 1 score</u>
Very good	0
Fairly good	1
Fairly bad	2
Very bad	3

Component 1 score: _____

Component 2: Sleep latency—questions 2 and 5a

<u>Response to Q2</u>	<u>Component 2/Q2 subscore</u>
≤ 15 minutes	0
16-30 minutes	1
31-60 minutes	2
> 60 minutes	3

<u>Response to Q5a</u>	<u>Component 2/Q5a subscore</u>
Not during past month	0
Less than once a week	1
Once or twice a week	2
Three or more times a week	3

<u>Sum of Q2 and Q5a subscores</u>	<u>Component 2 score</u>
0	0
1-2	1
3-4	2
5-6	3

Component 2 score: _____

Component 3: Sleep duration—question 4

<u>Response to Q4</u>	<u>Component 3 score</u>
> 7 hours	0
6-7 hours	1
5-6 hours	2
< 5 hours	3

Component 3 score: _____

Component 4: Sleep efficiency—questions 1, 3, and 4

Sleep efficiency = (# hours slept/# hours in bed) X 100%

hours slept—question 4

hours in bed—calculated from responses to questions 1 and 3

<u>Sleep efficiency</u>	<u>Component 4 score</u>
> 85%	0
75-84%	1
65-74%	2
< 65%	3

Component 4 score: _____

Component 5: Sleep disturbance—questions 5b-5j

Questions 5b to 5j should be scored as follows:

Not during past month	0
Less than once a week	1
Once or twice a week	2
Three or more times a week	3

<u>Sum of 5b to 5j scores</u>	<u>Component 5 score</u>
0	0
1-9	1
10-18	2
19-27	3

Component 5 score: _____

Component 6: Use of sleep medication—question 6

<u>Response to Q6</u>	<u>Component 6 score</u>
Not during past month	0
Less than once a week	1
Once or twice a week	2
Three or more times a week	3

Component 6 score: _____

Component 7: Daytime dysfunction—questions 7 and 8

<u>Response to Q7</u>	<u>Component 7/Q7 subscore</u>
Not during past month	0
Less than once a week	1
Once or twice a week	2
Three or more times a week	3

<u>Response to Q8</u>	<u>Component 7/Q8 subscore</u>
No problem at all	0
Only a very slight problem	1
Somewhat of a problem	2
A very big problem	3

<u>Sum of Q7 and Q8 subscores</u>	<u>Component 7 score</u>
0	0
1-2	1
3-4	2
5-6	3

Component 7 score: _____

Global PSQI Score: Sum of seven component scores: _____

Copyright notice: The Pittsburgh Sleep Quality Index (PSQI) is copyrighted by Daniel J. Buysse, M.D. Permission has been granted to reproduce the scale on this website for clinicians to use in their practice and for researchers to use in non-industry studies. For other uses of the scale, the owner of the copyright should be contacted.

Citation: Buysse, DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ: The Pittsburgh Sleep Quality Index (PSQI): A new instrument for psychiatric research and practice. *Psychiatry Research* 28:193-213, 1989