

Genetic characterization of *Cryptosporidium* spp. among patients with gastrointestinal complaints

Reza Ranjbar¹, Kaveh Baghaei², Ehsan Nazemalhosseini Mojarad²

¹Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

²Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

ABSTRACT

Aim: This study investigated subtypes of *Cryptosporidium* in patients with gastrointestinal complaints in Tehran, Iran.

Background: *Cryptosporidium*, an intracellular protozoan parasite, is among the major causative agents of gastroenteritis disorders in humans. It also causes water-borne and food-borne outbreaks of diarrheal diseases.

Patients and methods: A total of 1685 fecal samples were collected from patients with gastrointestinal complaints who had been referred to clinical laboratories Tehran, Iran. The primary diagnosis was established by the detection of oocysts using the modified Ziehl-Neelsen staining method and following that, the positive microscopically samples were selected for sequence analysis of the partial 60 kDa glycoprotein (gp60) gene.

Results: Out of 1685 collected samples, 7 (0.4 %) were positive for *Cryptosporidium* oocysts. Sequence analysis of gp60 gene in seven *Cryptosporidium* isolates revealed that two subtype families were identified, IIa and IIc. Five (of 7) isolates belonged to the subtype family IIa and the remaining two isolates belonged to IIc. Two sub-types were recognized within the subtype family IIa including IIaA16G2R1 (3/5), IIaA17G1R1 (2/5), while IIcA17G1d was the only subtype within IIc subtype family.

Conclusion: The predominance of zoonotic subtype families of *C. parvum* species (IIa, IIc) in this study highlights the importance of zoonotic transmission of cryptosporidiosis in the country.

Keywords: Genetic characterization, *Cryptosporidium*, Gastrointestinal complaints.

(Please cite as: Ranjbar R, Baghaei K, Nazemalhosseini Mojarad E. Genetic characterization of *Cryptosporidium* spp. among patients with gastrointestinal complaints referred to clinical laboratories of Tehran, Iran. *Gastroenterol Hepatol Bed Bench* 2016;9(4):301-307).

Introduction

Intestinal protozoan parasites are still major health problems in tropical and subtropical areas and are characteristically found among people with a low socio-economic grade and poor hygiene (1). Among these intestinal parasites,

Cryptosporidium spp. is a main pathogen which is a cause of diarrhea in children and immunocompromised patients and also infects the gastrointestinal tract of a wide range of vertebrates, including domestic and wild animals and birds (2-4).

The parasite is transmitted through the faecal-oral route, following direct or indirect contact with *Cryptosporidium* oocysts via person-to-person,

Received: 19 May 2016 Accepted: 18 August 2016

Reprint or Correspondence: Kaveh Baghaei, PhD. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran.

E-mail: kavehbaghai@gmail.com

zoonotic, contaminated water, foodborne or airborne contact (5-7).

Currently, the use of molecular approaches in genetic characterization of *Cryptosporidium* spp. at polymorphic loci has allowed a better understanding of the epidemiology of cryptosporidiosis (2-3). Molecular biology has well-known as a powerful tool for categorizing *Cryptosporidium* and has discovered major variation within the genus (about 30 species). Two predominantly species that have been found in humans are *C. parvum* and *C. hominis*. However, other species such as *C. meleagridis*, *C. muris*, *C. felis*, *C. canis*, *C. suis* and *C. andersoni* have been rarely detected in feces of immunocompetent and immunocompromised individuals (8).

In the past two decades, reports of cryptosporidiosis in Iran have been achieved using microscopy (9-13). Recently, researchers have developed PCR-based techniques for detection and identification of *Cryptosporidium* spp. (14-17).

Fingerprint of *C. parvum* infection has a critical role in outbreak investigations. DNA sequencing of the *Cryptosporidium* 60-kDa glycoprotein (GP60) gene has revealed substantial genetic heterogeneity among *C. hominis* and *C. parvum* isolates. GP60 gene could be used as a marker to determine the different subtype families within both species, including: Ia, Ib, Id, Ie, If and Ig for *C. hominis* and IIa, IIb, IIc, IId, IIe, II f , IIg, IIh, IIf, IIk, and III for *C. parvum* (18). Within each subtype group, there are several subtypes primarily based on the number of tri-nucleotide repeats coding for the amino acid serine (2, 7, 19).

To our knowledge, there are several molecular epidemiological studies that have documented the distribution of subtypes of *cryptosporidium* ssp. in children with diarrhea (8, 15), animals (20) and environments (the water that isolated from rivers) (21) in Iran.

In this study, we identified the genotypes of the *Cryptosporidium* isolates from patients with gastrointestinal complaints referred to clinical laboratories of Tehran using the polymerase chain reaction (PCR) amplification and sequencing analyses of the *Gp 60* gene.

Patients and Methods

Sampling

A total of 1685 fecal samples were collected from patients with gastrointestinal complaints who had been referred to Medical Centers in Tehran, Iran. *Cryptosporidium* oocysts were identified in samples after concentration by formalin–ethyl–acetate sedimentation and staining with a modified Zeihl-Neelsen technique (22). The positive *Cryptosporidium* spp. isolates were preserved in 2.5% potassium dichromate and kept at 4°C until DNA extraction.

DNA extraction

Approximately 300 µl of fecal suspension was washed three times with distilled water to remove trace of dichromate and then genomic DNA was extracted using DNAzol kit according to the manufacturer's instructions (Invitrogen, life technologies, Cat. No 10503-027, USA) with the addition of three freeze-thaw cycles (10 minutes) after resuspending samples in lysis buffer (to rupture the *Cryptosporidium* oocysts). The oocysts were frozen in the liquid nitrogen tank. Thawing was carried out at 90° C in the water bath.

DNA subtyping and sequence analysis

For subtyping *C. parvum* and *C. hominis*, a fragment of about 400 bp of the gp60 gene was amplified by nested PCR with the primers 5_-ATA GTC TCC GCT GTA TTC-3_ and 5_-GCA GAG GAA CCAGCA TC-3_ in primary PCR and 5_-TCC GCT GTA TTC TCA GCC-3_ and 5_-GAG ATA TAT CTT GGT GCG-3_ in secondary PCR, as described (Abe et al., 2006).

PCR products were visualized by electrophoresis in 1.5% agarose gels stained with ethidium bromide.

The PCR-amplified products were subjected to direct sequencing using a BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, Foster City, CA) and a Genetic Analyzer Prism™ 3130x1 (Applied Biosystems). The secondary PCR products were sequenced in both directions and, if variations were found, results were confirmed by sequencing of at least two independent PCR products. All sequences were edited manually and analyzed with reference sequences using the GenRunner software (v. 3.05).

Subtypes were recognized based on the number of trinucleotide repeats (TCA or TCG) coding for the amino acid serine (19).

Statistical analysis

The prevalence of *Cryptosporidium* infection and prevalence of *C. parvum* and *C. andersoni* in pre-weaned calves was compared with prevalence data for post-weaned calves. The Chi-square test for independence was used to analyze the data and differences were considered significant when $P < 0.05$. Statistical analysis was performed using SPSS (Ver. 12).

Results

Among the 1685 patients (66.3% male and 33.7% female) included in this study, microscopic examinations of the specimens revealed the presence of *Cryptosporidium*

oocysts in 7 (0.4 %) of the samples. Clinical information about these samples is presented in table 1. Patient complaints included abdominal pain, flatulence, tenesmus, diarrhea and dysentery. All samples were successfully amplified using specific primers (Figure 1) and PCR products of GP60 gene were purified and sequenced using a genetic analyzer machine.




Figure 1. PCR of *Cryptosporidium* based on GP60 gene. Lane 1: 100 bp DNA marker, lane A-D: *Cryptosporidium* spp.

The sequences were determined and analyzed using the chromas program and aligned with each other and with previously reported sequences for identification of the alleles and subtypes. The result of this analysis showed that all isolates were *C. parvum* species.

The sequence analysis of gp60 gene in seven *Cryptosporidium* isolates revealed that, two subtype families were identified, Ila and IId. Five (of 7) isolates belonged to the subtype family Ila and remaining two isolates belonged to IId. Two sub-types were recognized within the subtype family Ila including IlaA16G2R1 (3/5), and

Table 1. Distribution of *Cryptosporidium parvum* subtypes in isolates from patients with gastrointestinal complaints

ID	Subtype Family	Subtype	Sex/Age	GI complaints	Location
1	Ila	IlaA16G2R1	M\ 49	abdominal pain, tenesmus	Tehran
2	Ila	IlaA16G2R1	F\13	dysentery , abdominal pain	Varamin
3	Ila	IlaA16G2R1	M\ 32	abdominal pain	Pakdasht
4	Ila	IlaA17G1R1	M\27	dysentery , abdominal pain, flatulence	Tehran
5	Ila	IlaA17G1R1	F\ 52	abdominal pain	Pakdasht
6	IId	IIdA17G1d	M\11	dysentery , abdominal pain	Varamin
7	IId	IIdA17G1d	F\68	dysentery , abdominal pain, flatulence	Pakdasht

IlaA17G1R1 (2/5), while IIdA17G1d was the only subtype within the IId subtype family.

Discussion

There are wide intraspecific variations in *C. parvum* populations. Currently, at least 14 *C. parvum* subtype families, including Ila, I Ib, I Ic, I Id, I Ie, I If, I Ig, I Ih, I Ii, I Ik, I Il, I Im, I In, and I Io, have been identified on the basis of sequence analysis of the 60 kDa glykoprotein (GP60) gene (23).

In this study, *Cryptosporidium* isolated were classified by sequence analysis of the 60-kDa glycoprotein (GP60) gene. A prevalence rate of 0.4 % (7/1685) was obtained for cryptosporidiosis among patients with gastrointestinal complaints.

All *Cryptosporidium* isolates from patients with gastrointestinal complaints were identified as *C. parvum* species and none of them belonged to *C. hominis*. Also, two main subtype families (Ila and IId) were recognized. Regarding to other studies, *C. parvum* is reported in most cryptosporidiosis cases in Iran (6,14,24), which highlights the significance of zoonotic transmission of cryptosporidiosis in the country (25-26).

Although, in some countries such as Australia, India, Egypt, Mexico and Peru, predominance of *C. hominis* in human isolates have been documented (27-31), but our result is consistent with studies from some other countries such as Malaysia, Kuwait, Yemen, Sweden, United Kingdom, Netherland, France, Portugal, and Nicaragua (32-39)

First characterization of *Cryptosporidium* subtypes in Iranian specimens was documented in 2011. According to their results, 47 samples of *C. parvum* and *C. hominis* were characterized in children and cattle by sequence analysis of the gp60 gene, which showed cattle and children were mainly infected by *C. parvum* Ila subtypes and *C. parvum* Ila and IId subtypes, respectively (17). In some countries such as Spain, Egypt and China, IId subtypes are known to be more prevalent in

sheep and goats (40-42). Also, IId subtypes are also common in children from Iran neighboring countries (43-46). Sharbatkhori and her colleagues, showed three haplotypes of Ila subtype family including IlaA16G2R1, IlaA17G1R1, IlaA22G3R1 and one haplotype of IId subtype family among diarrheic children from Gonbad Kavoods City (Golestan Province, Northern Iran) and suggested a zoonotic transmission of cryptosporidiosis in this area (8).

The majority of Ila and IId subtypes highlight the significance of zoonotic *Cryptosporidium* transmission in Iran. Thus, cattle could be a plausible source of human infection with *C. parvum* Ila in Iran (15-18).

In another study, in 2011, high diversity of *Cryptosporidium* sub-genotypes was shown among Malaysian HIV infected individuals. The results obtained from this paper signified the possibility of zoonotic as well as anthroponotic transmissions of cryptosporidiosis in HIV infected individuals (47). In 2015, Mahmoudi, et al. determined the genotype and subtype distribution of *Cryptosporidium* spp. in river water samples in Iran. They showed that all *C. parvum* and *C. hominis* isolates belonged to the IId and Id subtype families, respectively and this source is a potential risk of waterborne cryptosporidiosis in humans and animals (21).

Vieira, et al. identified two subtype families (Ila and IId) from children, calves and eight pigs in Romania. They proposed, cattle might be the source of *Cryptosporidium* infections for humans and the transmission dynamics of *C. parvum* in Romania (48).

In their study, Wang, et al. suggest that, due to the higher nucleotide diversity of *C. parvum* IId GP60 sequences, more population genetic studies using high-resolution tools are needed to present a better explanation of the origin and dissemination of *C. parvum* in the world (23).

In 2010, sequence analysis of the GP60 locus identified three *C. parvum* and two *C. hominis*

subtype families in Jordan. In this study several rare and novel subtypes were reported as well (45).

Preliminary molecular epidemiological studies have revealed some unique features of cryptosporidiosis transmission in humans in Iran and other Mideast countries. As the *C. parvum* subtype family IId was the dominant family causing cryptosporidiosis in humans in Iran, zoonotic transmission could possibly be involved. However, more extensive sampling of both humans and farm animals, especially sheep and goats, and collection of epidemiological data in case-control and longitudinal studies are needed for a better understanding of the sources of *C. parvum* infections in humans in Iran and other Mideast countries.

References

- Haghighi A, Khorashad AS, Nazemalhosseini Mojarad E, Kazemi B, Rostami Nejad M, Rasti S. Frequency of enteric protozoan parasites among patients with gastrointestinal complaints in medical centers of Zahedan, Iran. *Trans R Soc Trop Med Hyg* 2009; 103: 452-54.
- Xiao L, Bern C, Sulaiman IM, Lal AA. Molecular epidemiology of human cryptosporidiosis. In: Thompson M, ed. *Cryptosporidium: from Molecules to Disease*. Amsterdam: Elsevier; 2004: 121-46.
- Xiao L. Molecular epidemiology of cryptosporidiosis: an update. *Exp Parasitol* 2010; 124: 80-9.
- Nazemalhosseini-Mojarad E, Keshavarz A, Taghipour N, Haghighi A, Kazemi B, Athari A. Genotyping of *Cryptosporidium* spp. in clinical samples: PCR-RFLP analysis of the TRAP-C2 gene. *Gastroenterol Hepatol Bed Bench* 2011; 4: 29-33.
- Fayer R. *Cryptosporidium*: a waterborne zoonotic parasite. *Vet Parasitol* 2004; 126: 37-56.
- Keshavarz A, Athari A, Haghighi A, Kazami B, Abadi A, Nazemalhosseini-Mojarad E, et al. Genetic characterization of *Cryptosporidium* spp. among children with diarrhea in Tehran and Qazvin Provinces, Iran. *Iranian J Parasitol* 2008; 3: 30-6.
- Plutzer J, Karanis P. Genetic polymorphism in *Cryptosporidium* species: an update. *Vet Parasitol* 2009; 165: 187-99.
- Sharbatkhori M, Nazemalhosseini Mojarad E, Taghipour N, Pagheh AS, Mesgarian F. Prevalence and genetic characterization of *Cryptosporidium* Spp. in diarrheic children from Gonbad Kavoos City, Iran. *Iran J Parasitol* 2015; 10: 441-47.
- Moghadam AA. Symptomatic and asymptomatic cryptosporidiosis in young children in Iran. *Pak J boil Sci* 2008; 10: 1108-12.
- Mirzaei M. prevalence of *Cryptosporidium* spp. infection in diarrheic and non diarrheic human in Iran. *Korean J Parasitol* 2007; 45: 133-37.
- Khalili B, Mardani M. Frequency of *Cryptosporidium* and risk factors related to cryptosporidiosis in under 5-year-old hospitalized children due to diarrhea. *Iranian J Clil Infec Dis* 2009; 4: 151-55.
- Hoghooghi-Rad N. Some epidemiological aspects of cryptosporidiosis in Ahvaz, center of Khoozestan Province, Islamic Republic of Iran. *Med J Islam Repub Iran* 1994; 1: 17-22.
- Nouri M, Moghadam A, Haghghatnia H. *Cryptosporidium* infection in human diarrhea patients in West Azerbaijan, Iran. *Med J Islam Repub Iran* 1991; 2: 35-8.
- Meamar AR, Rezaian M, Rezaie S, Mohraz M, Mohebbali M, Mohammad K, et al. SSU-rRNA gene analysis of *Cryptosporidium* spp. in HIV positive and negative patients. *Iranian J Publ Health* 2006; 35: 1-7.
- Taghipour N, Nazemalhosseini-Mojarad E, Haghighi A, Rostami-Nejad M, Romani S, Keshavarz A, et al. Molecular epidemiology of *Cryptosporidiosis* in Iranian children admitted to a pediatric hospital in Tehran, Iran. *Iranian J Parasitol* 2011; 6: 41-5.
- Nazemalhosseini-Mojarad E, Taghipour N, Haghighi A, Keshavarz A, Rostami Nejad M, Zali MR. DNA fingerprinting of bovine *Cryptosporidium* isolates in Qazvin Province. *Iran Koomesh* 2011; 12: 408-12.
- Nazemalhosseini-Mojarad E, Haghighi A, Taghipour N, Keshavarz A, Mohebbi SR, Zali MR, et al. Subtype analysis of *Cryptosporidium parvum* and *C. hominis* isolates from humans and cattle in Iran. *Vet Parasitol* 2011; 179: 250-52.
- Nazemalhosseini-Mojarad E, Feng Y, Xiao L. The importance of subtype analysis of *Cryptosporidium* spp. in epidemiological investigations of human cryptosporidiosis in Iran and other Mideast countries. *Gastroenterol Hepatol Bed Bench* 2012; 5: 67-70.

19. Sulaiman IM, Hira PR, Zhou L, Al-Ali FM, Al-Shelahi FA, Shweiki HM, et al. Unique endemicity of Cryptosporidiosis in children in Kuwait. *J Clin Microbiol* 2005; 43: 2805-809.
20. Keshavarz A, Haghighi A, Athari A, Kazemi B, Abadi A, Nazemalhosseini-Mojarad E. Prevalence and molecular characterization of bovine *Cryptosporidium* in Qazvin Province, Iran. *Vet Parasitol* 2009; 160: 316-18.
21. Mahmoudi MR, Nazemalhosseini-Mojarad E, Kazemi B, Haghighi A, Mirzaei A, Mohammadiha A, et al. *Cryptosporidium* genotypes and subtypes distribution in river water in Iran. *J Water Health* 2015; 13: 600-6.
22. Casemore DP, Armstrong M, Sands RLLaboratory diagnosis of cryptosporidiosis. *J Clin Pathol.* 1985; 38: 1337-41.
23. Wang R, Zhang L, Axén C, Bjorkman C, Jian F, Amer S, et al. *Cryptosporidium parvum* IId family: clonal population and dispersal from Western Asia to other geographical regions. *Sci Rep* 2014; 4: 4208.
24. Pirestani M, Sadraei J, Dalimi asl A, Zavvar M, Vaeznia H. Molecular characterization of *Cryptosporidium* isolates from human and bovine using 18s rRNA gene in Shahriar county of Tehran, Iran. *Parasitol Res* 2008; 103: 467-72.
25. Izadi M, Jonaidi-Jafari N, Saburi A, Eyni H, Rezaeiamesh MR, Ranjbar R. Cryptosporidiosis in Iranian farm workers and their household members: a hypothesis about possible zoonotic transmission. *J Trop Med* 2014; 2014: 405875.
26. Izadi M, Jonaidi-Jafari N, Saburi A, Eyni H, Rezaeiamesh MR, Ranjbar R. Prevalence, molecular characteristics and risk factors for Cryptosporidiosis among Iranian immunocompromised patients. *Microbiol Immunol* 2012; 56: 836-42.
27. Cama VA, Bern C, Roberts J, Cabrera L, Sterling CR, Ortega Y, Gilman RH, Xiao L. *Cryptosporidium* species and subtypes and clinical manifestations in children. *Peru Emerg Infect Dis* 2008; 14: 1567-74.
28. Valenzuela O, González-Díaz M, Garibay-Escobar A, Burgara-Estrella A, Cano M, Durazo M, et al. Molecular characterization of *Cryptosporidium* spp. in children from Mexico. *PLoS ONE* 2014; 9: e96128.
29. Sharma P, Sharma A, Sehgal R, Malla N, Khurana S. Genetic diversity of *Cryptosporidium* isolates from patients in North India. *Int J Infect Dis* 2013; 17: e601-e05.
30. Helmy YA, Krücken J, Nöckler K, von Samson-Himmelstjerna G, Zessin KH. Molecular epidemiology of *Cryptosporidium* in livestock animals and humans in the Ismailia province of Egypt. *Vet Parasitol* 2013; 193: 15-24.
31. O'Brien E, McInnes L, Ryan U. *Cryptosporidium* GP60 genotypes from humans and domesticated animals in Australia, North America and Europe. *Exp Parasitol* 2008; 118: 118-21.
32. Alyousefi NA, Mahdy MA, Lim YA, Xiao L, Mahmud R. First molecular characterization of *Cryptosporidium* in Yemen. *Parasitology* 2013; 140: 729-34.
33. Guyot K, Follet-Dumoulin A, Lelievre E, Sarfati C, Rabodonirina M, Nevez G, et al. Molecular characterization of *Cryptosporidium* isolates obtained from humans in France. *J Clin Microbiol.* 2001; 39: 472-80.
34. Lim YA, Iqbal A, Surin J, Sim BL, Jex AR, Nolan MJ, et al. First genetic classification of *Cryptosporidium* and *Giardia* from HIV/AIDS patients in Malaysia. *Infect Genet Evol* 2011; 11: 968-74.
35. Alves M, Xiao L, Sulaiman I, Lal AA, Matos O, Antunes F. Subgenotype analysis of *Cryptosporidium* isolates from humans, cattle, and zoo ruminants in Portugal. *J Clin Microbiol* 2003; 41: 2744-47.
36. Muñoz-Antoli C, Pavón A, Marcilla A, Toledo R, Esteban JG. Prevalence and molecular characterization of *Cryptosporidium* in schoolchildren from department of Rio San Juan (Nicaragua). *Trop Biomed* 2011; 28: 40-7.
37. Wielinga PR, de Vries A, van der Goot TH, Mank T, Mars MH, Kortbeek LM, et al. Molecular epidemiology of *Cryptosporidium* in humans and cattle in The Netherlands. *Int J Parasitol* 2008; 38: 809-17.
38. Insulander M, Silverlås C, Lebbad M, Karlsson L, Mattsson JG, Svenungsson B. Molecular epidemiology and clinical manifestations of human Cryptosporidiosis in Sweden. *Epidemiol Infect* 2013; 141: 1009-20.
39. McLauchlin J, Amar C, Pedraza-Díaz S, Nichols GL. Molecular epidemiological analysis of *Cryptosporidium* spp. in the United Kingdom: results of genotyping *Cryptosporidium* spp. in 1,705 fecal samples from humans and 105 fecal samples from livestock animals. *J Clin Microbiol* 2000; 38: 3984-90.
40. Quilez J, Torres E, Chalmers RM, Hadfield SJ, Del Cacho E, Sanchez-Acedo C. Genotype and subtype characterization of *Cryptosporidium* in lambs and goat kids in Spain. *Appl Environ Microbiol* 2008; 74: 6026-31.
41. Amer S, Honma H, Ikarashi M, Tada C, Fukuda Y, Suyama Y, et al. *Cryptosporidium* genotypes and

- subtypes in dairy calves in Egypt. *Vet Parasitol* 2010; 169: 382-86.
42. Wang R, Wang H, Sun Y, Zhang L, Jian F, Qi M, et al. Characteristics of *Cryptosporidium* transmission in pre-weaned dairy cattle in Henan, China. *J Clin Microbiol* 2011; 49: 1077-82.
43. Tamer GS, Turk M, Dagci H, Pektas B, Guy EC, Guruz AY, et al. The prevalence of Cryptosporidiosis in Turkish children, and genotyping of isolates by nested polymerase chain reaction-restriction fragment length polymorphism. *Saudi Med J* 2007; 28: 1243-46.
44. Al-Brikan FA, Salem HS, Beeching N, Hilal N. Multilocus genetic analysis of *Cryptosporidium* isolates from Saudi Arabia. *J Egypt Soc Parasitol* 2008; 38: 645-58.
45. Hijjawi N, Ng J, Yang R, Atoum MF, Ryan U. Identification of rare and novel *Cryptosporidium* GP60 subtypes in human isolates from Jordan. *Exp Parasitol* 2010; 125: 161-64.
46. Iqbal J, Khalid N, Hira PR. Cryptosporidiosis in Kuwaiti children: association of clinical characteristics with *Cryptosporidium* species and subtypes. *J Med Microbiol* 2011; 60: 647-52.
47. Iqbal A, Lim YAL, Surin J, Sim BLH. High Diversity of *Cryptosporidium* subgenotypes identified in Malaysian HIV/AIDS individuals targeting gp60 gene. *PLoS ONE* 2012; 7: e31139.
48. Vieira PM, Mederle N, Lobo ML, Imre K, Mederle O, Xiao L, Darabus G, Matos O. Molecular characterisation of *Cryptosporidium* (Apicomplexa) in children and cattle in Romania. *Folia Parasitol (Praha)* 2015; 62. pii: 2015.002