

Farida Abesi¹ DDS, MS, Maryam Ehsani^{2*} DDS, MS

Radiographic evaluation of maxillary anterior teeth canal curvatures in an Iranian population

1. Assistant Professor of Oral and Maxillofacial Radiology, Dental school, Babol University of Medical Sciences, Babol, Iran.

2. Assistant Professor of Endodontics. Dental school, Babol University of Medical Sciences, Babol, Iran. *Correspondence author Email: ehsanimaryam@yahoo.com

INTRODUCTION: Complete knowledge of root canal curvature is a critical factor in successful endodontic treatment. The aim of this study was to investigate the direction, radius and degree of curvature of maxillary anterior teeth and the relationship between the radius and degree of curvature in Babol, a northern city of Iran.

MATERIALS & METHODS: A total of 242 radiographs of maxillary anterior teeth (central, lateral and canine) were taken by periapical parallel technique and processed by automatic processing. The degree of canal curvature was measured only at mesiodistal direction with Schneider method and classified according to Seidberg method. Statistical analysis was performed with Kruskal Wallis and Mann-Whitney *U* tests.

RESULTS: Overall, 153 (62%) teeth had curvatures; 35.3% were mesially inclined and 64.7% were distally inclined. The degree of canal curvature was categorized into small, intermediate, and severe, that is 39.3%, 44.6% and 16.1%, respectively. The mean value of root curvature angle was $7.24 \pm 9.03^\circ$ in central incisors, $12.08 \pm 11.02^\circ$ in lateral incisors, and $15.08 \pm 12.02^\circ$ in canines respectively. There was significant correlation between type of tooth and degree of curvature ($P=0.000$). Significant correlation was not found between the type of tooth and radius of curvature ($P=0.365$).

CONCLUSION: In the present study, 62% of maxillary anterior teeth had some form of curvatures; highest degrees of curvature were attributed to the canine teeth.

KEYWORDS: Maxillary anterior teeth, Radius of curvature, Root curvature.

Received Oct 2010; accepted Dec 2010

INTRODUCTION

The main object of root canal therapy is the elimination of microorganisms and infected tissue from the tooth root canal system. This is performed by enlarging and shaping the canals to allow for adequate chemical debridement, while at the same time preserving the original shape and structure of the tooth (1). Thorough knowledge of the anatomical configurations of the dental pulp, and the possible variations is critical for successful endodontics (2). Conducting root canal therapy (RCT) when uncertain of the canal morphology increases the risk of transportation, ledge formation and even perforation, and often results in failure of the root canal procedure (3-5). Several techniques are used to determine root canal configuration

such as specimen transparent technique, conventional radiographs, radiopaque contrast media, cross-sectional cutting scanning electron microscopy (SEM) and cone beam computed tomography (CBCT) (6-11).

Schneider divided the root canal curvature into different root angles (12). However, evaluating the curvature by the angle of the canal is now thought to be insufficient. For instance, two canals with the same angle obtained by the Schneider method could have very different radii of curvature or abruptness of curvatures (13). Pruett *et al.* proposed that the assessment of canal curvature should be indicated by two measurements: *a)* the angle of curvature and *b)* the radius of curvature determined mathematically




Figure 1. Angel (α) and Radius (r) of root curvature

from radiographs (14). There is a general shortage of clinical studies on canal curvature of maxillary anteriors as most investigations have concentrated on the type and variation of the root canal system (8,15). Few investigations have been conducted on the specific degree of canals curvature of the maxillary anterior teeth (16). The purpose of this study was to determine the degree, direction and radius of canal curvature in maxillary anterior teeth in Babol, a northern city of Iran.

MATERIALS & METHODS

This study was performed on 242 radiographs of anterior teeth (central, lateral, and canine) taken from patients referred to a maxillofacial radiology clinic in Babol, Iran. The patients were initially informed about the study and upon agreement they filled the inform consent form. Teeth that had not received endodontic therapy and had complete apical foramen and contour were selected. All radiographs were taken by periapical parallel technique with film holder XCP (Dentsply, United Kingdom) and processed by automatic processor (Velopex, United Kingdom).

The angle of curvature was determined by the Schneider technique which measures curvature as well as the acute angle between the long axis of the canal and a line joining the apical foramen to the point of initial curvature shown on the radiograph (Figure 1) (12). Teeth were classified according to Seidberg classification; the degree of the curvature was categorized as low ($<5^\circ$), moderate ($5-25^\circ$) and severe ($25-70^\circ$). The radius of curvature was measured by the Estrella method; that is, two 6-mm semi straight lines superimposed to root canal (Figure 1) (11).

According to this method, the first line (line **b**) represents longer continuity of the apical region and the second line (line **a**) represents the middle and cervical thirds. Of the second line, the nearest 6mm to the first line was considered; the lines perpendicular to the midpoints of two 6-mm semi straight lines convene each other at the point that is Circum Center. This point is the center of the circle which determines the magnitude of the root curve (11). The curvature measurement is performed by two observers with conveyor (Rotting, Germany) with accuracy of one degree. According the Estrella, the root curvature radius was classified as following:

- Small radius ($\leq 4\text{mm}$) *i.e.* severe curvature
- Intermediate radius ($4 < r \leq 8$) *i.e.* moderate curvature
- Large radius ($r > 8\text{mm}$) *i.e.* mild curvature

The data were analyzed with Kruskal Wallis and Mann-Whitney *U* tests. For multiple comparisons bonferroni adjustment of α was carried out.

RESULTS

Out of 242 examined teeth, 153 (62%) had curvatures; 35.3% were mesially and 64.7% were distally inclined (Table 1).

The degree of canal curvature was as follows: 39.3% with small curvature, 44.6% with intermediate and 16.1% with sever curvature (Table 2).

Distribution of different curvature radius (according to Estrella) in studied teeth is presented in Table 3.

The tooth type showed no significant correlation with radius of curvature ($P=0.365$) (Table 4), and significant correlation with curvature angle ($P=0.000$) (Table 5). Mann-Whitney *U* test shown significant difference between central and lateral ($P=0.003$), central and canine ($P=0.000$) but the difference between lateral and canine was not significant ($P=0.128$).

DISCUSSION

Radiographs are commonly utilized in endodontics to determine canal morphology;

Table 1. Frequency of mesial+distal curvatures among the teeth

Direction	Mesial	Distal	Total
Frequency (%)	54 (35.3)	99 (64.7)	153 (100.0)

Table 2. Frequency and percentage of angles of curvatures among the teeth

Angle	≤4	5-24	≥25	Total
Frequency (%)	95(39.3)	108 (44.6)	39 (16.1)	242 (100.0)

Table 3. Frequency and percentage of radius in teeth with angles over 20

Radius	≤4	5-8	≥9	Total
Frequency (%)	2 (2.5)	15 (19.0)	62 (78.5)	79 (100.0)

Table 4. Mean and standard deviation of radius according to tooth type

Tooth	Number	Mean±SD	P value
Central	18	12.44±3.36	0.365
Lateral	35	12.00±5.00	
Canine	26	10.84±3.22	

Table 5. Mean and standard deviation of degree of curvature according to tooth type

Tooth	Number	Mean±SD	P value
Central	86	7.24±9.03	0.000
Lateral	100	12.08±11.02	
Canine	56	15.08±12.02	

however the two dimensional image of the teeth might cause some uncertainty as to the actual direction and the degree of root canal curvature. Radiographs are essential to the practice of endodontics (17); when assessing root canal curvatures parallel technique is the method of choice as it provides a less distorted view of the dentition, as well as clearer apical areas. Bisecting angle technique may be used if there are anatomic constraints such as a shallow palate. However, in this study these cases were excluded. Endodontic cases were also excluded, as K-files do not exactly conform to the canal shape, and will not always remain centered in the canal lumen, moreover iatrogenic errors may occur (18,19). Several studies (14,20-26) have suggested methods to determine root canal curvature using periapical radiographs. Our technique determined angle of root curvature based on Schneider method (12).

In present study, the degree of root canal curvature of maxillary canines was larger than that of maxillary incisors concurring with the findings of Tao *et al.* They investigated root canal curvature in 400 maxillary anterior teeth by indirect digital radiography from both labiolingual and mesiodistal direction. He also concluded that the radius of curvature of maxillary canines root canals were smaller than that of maxillary incisors (16). However, our study was not able to find a significant relation between type of tooth and the radius curvature (P=0.365).

Estrella *et al.* described a new method to determine root curvature radius in CBCT images (12). According to this method we measured the radius of root curvature and found that there was inverted relation between radius and degree of root curvature; however this was not significant ($r=0.155$, $P>0.05$).

Our results agree with two previous studies that found no significant difference between the degree and the radius of curvature of root canals; also no relationship existed between the original radius of curvature and apical transportation (14,20). However, determination of root canal curvature by use of its radius has proven to be an effective method.

Further studies should concentrate on CBCT images of teeth in buccolingual and mesiodistal directions, keeping dose limitation in mind.

CONCLUSION

In the present study, 62% of maxillary anterior teeth had root curvatures and the lowest degree of curvature was related to the centrals. The results of this study can enhance endodontic therapy predictability and minimize errors during post insertion.

Conflict of Interest: 'none declared'.

REFERENCES

- Schilder H. Cleaning and shaping the root canal. *Dent Clin North Am* 1974;18:269-96.
- Rahimi S, Shahi S, Lotfi M, Zand V, Abdolrahimi

- M, Es'haghi R. Root canal configuration and the prevalence of C-shaped canals in mandibular second molars in an Iranian population. *J Oral Sci* 2008;50:9-13.
3. Mullaney TP. Instrumentation of finely curved canals. *Dent Clin North Am* 1979;23:575-92.
 4. Weine FS, Kelly RF, Bray KE. Effect of preparation with endodontic handpieces on original canal shape. *J Endod* 1976;2:298-303.
 5. Roane JB, Sabala CL, Duncanson MG Jr. The "balanced force" concept for instrumentation of curved canals. *J Endod* 1985;11:203-11.
 6. Miyashita M, Kasahara E, Yasuda E, Yamamoto A, Sekizawa T. Root canal system of the mandibular incisor. *J Endod* 1997;23:479-84.
 7. Kartal N, Cimilli HK. The degrees and configurations of mesial canal curvatures of mandibular first molars. *J Endod* 1997;23:358-62.
 8. Naoum HJ, Chandler NP, Love RM. Conventional versus storage phosphor-plate digital images to visualize the root canal system contrasted with a radiopaque medium. *J Endod* 2003;29:349-52.
 9. Lee MM, Rasimick BJ, Turner AM, Shah RP, Musikant BL, Deutsch AS. Morphological measurements of anatomic landmarks in pulp chambers of human anterior teeth. *J Endod* 2007;33:129-31.
 10. Gilles J, Reader A. An SEM investigation of the mesiolingual canal in human maxillary first and second molars. *Oral Surg Oral Med Oral Pathol* 1990;70:638-43.
 11. Estrela C, Bueno MR, Sousa-Neto MD, Pécora JD. Method for determination of root curvature radius using cone-beam computed tomography images. *Braz Dent J* 2008;19:114-8.
 12. Schneider SW. A comparison of canal preparations in straight and curved root canals. *Oral Surg Oral Med Oral Pathol* 1971;32:271-5.
 13. Zheng QH, Zhou XD, Jiang Y, Sun TQ, Liu CX, Xue H, Huang DM. Radiographic investigation of frequency and degree of canal curvatures in Chinese mandibular permanent incisors. *J Endod* 2009;35:175-8.
 14. Pruett JP, Clement DJ, Carnes DL Jr. Cyclic fatigue testing of nickel-titanium endodontic instruments. *J Endod* 1997;23:77-85.
 15. Sert S, Aslanalp V, Tanalp J. Investigation of the root canal configurations of mandibular permanent teeth in the Turkish population. *Int Endod J* 2004;37:494-9.
 16. Tao XL, Peng B, Bian Z, Fan MW. Survey of root canal curvature in maxillary anterior teeth. *Hua Xi Kou Qiang Yi Xue Za Zhi*. 2007;25:135-8.
 17. Intraoral radiographic examination. In: Michael Pharaoh: *Oral Radiology*, 6th Edition. St. Louis: CV Mosby, 2009: pp.150.
 18. Jerome CE, Hanlon RJ Jr. Identifying multiplanar root canal curvatures using stainless-steel instruments. *J Endod* 2003;29:356-8.
 19. Cunningham CJ, Senia ES. A three-dimensional study of canal curvatures in the mesial roots of mandibular molars. *J Endod* 1992;18:294-300.
 20. Lopes HP, Elias CN, Estrela C, Siqueira JF Jr. Assessment of the apical transportation of root canals using the method of the curvature radius. *Braz Dent J* 1998;9:39-45.
 21. Lopes HP, Moreira EJ, Elias CN, de Almeida RA, Neves MS. Cyclic fatigue of ProTaper instruments. *J Endod* 2007;33:55-7.
 22. Moreira EJJ, Lopes HP, Elias CN, Fidel RAS. Flexion rotation fracture of NiTi endodontic instruments. *RBO* 2002;59:412-4.
 23. Pecora JD, Capelli A, Guerisoli DM, Spanó JC, Estrela C. Influence of cervical preflaring on apical file size determination. *Int Endod J* 2005;38:430-5.
 24. Schäfer E, Florek H. Efficiency of rotary nickel-titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals. *Int Endod J* 2003;36:199-207.
 25. Nagy CD, Szabó J, Szabó J. A mathematically based classification of root canal curvatures on natural human teeth. *J Endod* 1995;21:557-60.
 26. Willershausen B, Kasaj A, Tekyatan H, Roehrig B, Briseno B. Radiographic investigation of location and angulation of curvatures in human maxillary incisors. *J Endod* 2008;34:1052-6.