

The role of socioeconomic welfare in the prevalence of severe vitamin D deficiency, vitamin B12 deficiency and glucose metabolic changes in population of Karaj, Iran

Mahsa Houshdar*¹, Seyed Mehdi Samimi Ardestani², Seyed Saeed Sadr³

¹. MD. CEO of Houshdar Medical Technology Co, Iran. (Corresponding Author: mhooshdar@gmail.com)

². M.D , Associate professor of psychiatry, Shahid Beheshti University of Medical Science, behavioral science research center, Imam Hossein Medical Center, Tehran, Iran

³. MD , assistant professor of psychiatry , Shahid Beheshti University Of Medical Science, behavioral science research center, Imam Hossein Medical Center, Tehran, Iran

(Received: 20 Oct 2015; Revised: 8 Nov 2015; Accepted: 26 Nov 2015)

Abstract

Introduction: Online Medicare is a method in which parts of a medical process, whether its diagnostics, monitoring or the treatment itself will be completed by using online services. At the first step the students were registered for using the system. They participated in estimating depression scale; anxiety scale and clinical interview by online medical care system. Subsequently, the lab examination tests were performed on persons specified by the system. The lab examinations include: serum level of vitamin D^{3&4}, serum level of vitamin B12^{5&6}, fasting blood sugar^{7&8}, HbA1c^{7&8}, thyroid function test^{9&10} and CBC. All of the students were solely treated by vitamins or minerals therapy and/or treatment of medical problems (such as hypothyroidism).

Methods: For detecting the role of socioeconomic welfare in the prevalence of the diseases, we implement the system in a low middle class boy's state high school, and in an above average boy's state high school and in an above average boy's private high school. The prevalence of severe vitamin D deficiency, hypoglycemia, impaired fasting glucose level and vitamin B12 deficiency were compared in students who lay in different socioeconomic situation.

Results: Severe vitamin D deficiency is significantly higher in above average neighborhoods than low middle class neighborhoods with p value = 0.024 < 0.05. Significant difference in vitamin B12 deficiency, hypoglycemia and impaired fasting glucose were seen in deferent high schools, but it is not related to socioeconomic situation.

Conclusion: In Karaj, Iran, in societies with better socioeconomic situation, we can find higher severe vitamin D deficiency, it can be due to urbanization, living in apartment, over protection, preferring indoor games to outdoor games.

Declaration of Interest: None.

Key words: Depression, Anxiety, Vitamin D Deficiency, Socioeconomic.

Introduction

Online Medicare is a method in which parts of a medical process, whether its diagnostics, monitoring or the treatment itself will be completed by using online services(1). For detecting the role of socioeconomic welfare in the prevalence of the diseases, we implement the system in a lower middle class boy's state high school, and in an

above average boy's state high school and in an above average boy's private high school.

Methods

At the first step the students were registered for using the system. It was not mandatory and not free. They participated in estimating depression scale; anxiety scale and clinical interview by online medical care system. During this estimation, we

could find out about the existence and severity of depression and anxiety in each one of the participants. Also we could determine the consequent needs of each one, such as supportive therapy¹ in mild depression or anxiety, need to be visited by psychologist in moderate cases(1), need to be visited by psychiatrist in moderate-severe cases¹, need to be visited by psychiatrist and psychologist in severe cases and need to perform medical lab examination tests (2). The lab examination tests were performed on persons specified by the system. The lab

examinations included: serum level of vitamin D(3-4), serum level of vitamin B12^{5&6}, fasting blood sugar(7-8), HbA1c(7-8), thyroid function tests(9-10) and CBC. All of the students were solely treated by vitamins or minerals therapy and/or treatment of medical problem (such as hypothyroidism).


Results

Descriptive findings related to variables including mean and standard deviation.

Table 1. The prevalence of severe vitamin D deficiency, vitamin B12 deficiency and glucose metabolism changes in different neighborhoods.

	Vit D<10	Vit B12 <200	Hypoglycemia	Impaired Fasting Glucose Level
Lower middle class boy's state high school	4 %10.52	2 %5.2	1 %2.6	8 %21
Above average boy's state high school	31 %41.3	20 %26	6 %8	1 %1.3
Above average boy's private high school	25 %44.64	1 %1.78	0 %0	8 %14.28

Chart 1. The prevalence of severe vitamin D deficiency, vitamin B12 deficiency and glucose metabolism changes in different neighborhoods.



Series 1- Low middle class boy's state high school
 Series 2- Above average boy's state high school
 Series 3- Above average boy's private high school
 Category 1-Vit D<10
 Category 2-Vit B12 <200
 Category 3-Hypoglycemia
 Category 4-Impaired Fasting Glucose Level

Statistical Results:

- Severe vitamin D deficiency is significantly higher in above average neighborhoods than lower middle class neighborhoods with p value = $0.024 < 0.05$.
 - Hypoglycemia is higher in the state high schools in above average neighborhood than state high schools in lower middle class neighborhood and private high school above average neighborhood.
 - Severe vitamin B12 deficiency is significantly higher in the state high schools in above average neighborhood than state high schools in lower middle class neighborhood and private high schools above average neighborhood.
- Impaired fasting glucose level is lower in the state high schools in above average neighborhood than state high schools in lower middle class neighborhood and private high schools above average neighborhood

Conclusion

Reasons of higher severe vitamin D deficiency in above average neighborhood than lower middle class neighborhood:

- Urbanization: Apartment living is very popular in above average neighborhoods. Most of these apartments have not enough outdoor space available, thus direct sunlight availability is rare¹¹.
- Over protection: In above average neighborhoods the students are not permitted to exit home alone. Transferring to school is mostly performed by private vehicle or school vehicle. Naturally they do not have enough direct sunlight exposure time¹¹.
- Change in habits: Computer games, computer and internet is primary entertainment of students in above average neighborhoods, outdoor games such as football is not as popular as a few years ago. Thus, there is not enough interest for exposure to sunlight¹¹.
- The other results cannot be interpreted by available information; the future studies can help interpreting this data.

Acknowledgements:

- Thanks to everyone who helped us and led us in performing this project, including:
- Dr. Hosain Ziaodiny, General Director of Health, Ministry of Education

- Reza Taremy, Head of Health, Department of Education, Alborz Province
- Dr. Tahery, Expert in Adolescent Health Department in Ministry of Health
- Fatemeh Aghazadeh, Expert in Health and Prevention of Social Problems, Ministry of Education, Alborz Province, Third District
- Majid Gholami, Head of Health and Prevention of Social Problems, Ministry of Education, Alborz province, Third District

Acknowledgment

We would like to thank to our colleagues and the organizations for all provided insight and expertise that greatly assisted this research and patients who helped us kindly in the project. We also tried to consider all ethical issues in this study.

References

1. Houshdar m. Online medicare. Yerevan, armenia: world psychiatric association thematic conference "mental health and mental illness: focusing on eurasia; august 29-31, 2013.p.
2. Houshdar M. Results of operating online medical care system in high schools. The United States of America: Journal of Psychology & Clinical Psychiatry, volume 2, Issue 3; 2015. P.
3. Faghih S1, Abdolazadeh M1, Mohammadi M1, Hasanzadeh J2 Int J Prev Med. Prevalence of vitamin d deficiency and its related factors among university students in shiraz, iran: Int J Prev Med; 2014 Jun.5(6):796-9.
4. Anglin RE1, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. St Joseph's Hospital, 50 Charlton Avenue E, Hamilton, Ontario, Canada: Br J Psychiatry; 2013.P.100-7
5. Beers MH, Poter RS, Jones TV, Kaplan JL, Berkwits M. The Merck Manual of Diagnosis and Therapy :vitamin B12 deficiency, symptoms & signs, 18th edition, USA: Merck Research Laboratories, 2006, P. 38-39.
6. Fakhrzadeh H1, Ghotbi S, Pourebrahim R, Nouri M, Heshmat R, Bandarian F, Shafae A, Larijani B. Total plasma homocysteine, folate, and vitamin B12 status in healthy Iranian adults: the Tehran homocysteine survey (2003–2004)/a cross – sectional population based study. Iran: BMC Public Health; 2006 Feb .P. 6-29.
7. Beers MH, Poter RS, Jones TV, Kaplan JL, Berkwits M. The Merck Manual of Diagnosis and Therapy: diabetes mellitus (DM), other complication , 18th edition, USA: Merck Research Laboratories, 2006.P.1668.

8. Amini M1, Janghorbani M. Rev Diabet Stud. Diabetes and impaired glucose regulation in first-degree relatives of patients with type 2 diabetes in isfahan, iran: prevalence and risk factors. Iran. Journal of the society for biomedical diabetes research. 2007 Fall;4(3). P.169-76.
9. Beers MH, Poter RS, Jones TV, Kaplan JL, Berkwits M. The Merck Manual of Diagnosis and Therapy: subclinical hypothyroidism , 18th edition, USA: Merck Research Laboratories, 2006.P.1202-1203.
10. Delshad H, Mehran L, Tohidi M, Assadi M. The incidence of thyroid function abnormalities and natural course of subclinical thyroid disorders, Tehran, I.R. Iran: J Endocrinol Invest: 2012 May;35(5). P.516-21
11. Wacker M1, Holick MF1. Sunlight and Vitamin D: A global perspective for health. Dermatoendocrinol. 2013 Jan 1;5(1). P.51-108