

Effectiveness of Neurogenesis in treating Children with Cerebral Palsy

How to Cite this Article: Amirsalary S, Dehghan L, Dalvand H, Haghgoo H. Effectiveness of Neurogenesis in treating children with Cerebral Palsy. Iran. J. Child Neurol. 2012;6(2):1-8.

Susan AMIRSALARI MD¹,
Leila DEGHAN ²,
Hamid DALVAND²,
Hojjat Allah HAGHGOO ³

1. Associate Professor, Department of Pediatric Neurology, Faculty of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran

2. PhD Student of Occupational Therapy, Department of Occupational Therapy, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

3. Assistant Professor, Department of Occupational Therapy, University of Social Welfare & Rehabilitation Sciences, Tehran, Iran

Corresponding Author:

Dehghan L.

Address: Department of Occupational Therapy, University of Social Welfare & Rehabilitation Sciences, Tehran, Iran

Tel: +98 21 22180037

Fax: +98 21 22180109

Email: le.dehghan@uswr.ac.ir

Received: 18-Dec- 2011

Last Revised: 28-May-2012

Accepted: 2-June-2012

Objective

Tissue-specific stem cells divide to generate different cell types for the purpose of tissue repair in the adult. The aim of this study was to detect the significance of neurogenesis in the central nervous system in patients with cerebral palsy (CP).

Materials & Methods

A search was made in Medline, CINAHL, PubMed, ISI Web of Science and Google Scholar from 1995 to February 2011. The outcomes measured in the review were classified to origins, proliferation, and migration of new neurons, and neurogenesis in CP.

Results

According to the review of articles, neurogenesis persists in specific brain regions throughout lifetime and can be enhanced from endogenous progenitor cells residing in the subventricular zone by growth factors or neurotrophic factors and rehabilitation program.

Conclusion

Most of the studies have been conducted in the laboratory and on animals, more work is required at the basic level of stem cell biology, in the development of human models, and finally in well-conceived clinical trials.

Keywords: Stem cells; Neurogenesis; Cerebral palsy

Introduction

Plasticity is derived from the Greek word *plaistikos* meaning to form. This word mentions the brain's proficiency in learning, remembering and forgetting, together with its replanning and injury recovery potential (1). Plasticity involves the change of excitation and inhibition stability; a long-term potentiation (LTP) or long term depression (LTD); alteration of the neuronal membrane excitability; and finally anatomical changes, mainly construction of new axon terminals and new synapses (2).

Plasticity is a characteristic of the young human brain. Disorders and brain lesions such as cerebral palsy (CP) that occur during the developmental process involve the natural course of the brain evolution (3). CP is a disorder that influences movement and posture development; subsequently leading to restriction of activity (4). CP has been reported in 2 to 2.6 of 1000 live births in Iran (5), which is reasonably similar to other countries (6,7). As a result of better neonatal care, the survival of premature infants is increasing; therefore, CP is seen more often (8). Periventricular leukomalacia (PVL) injury is the cause of spastic diplegia, which is the most common form of CP. This defect occurs before 30-32

weeks (9). An increasing regional gray matter volume due to axonal sprouting, neuronal hypertrophy and neurogenesis is reported in preterm children affected by PVL, which may show brain plasticity (10). Therefore, neurogenesis is one of the neuroplasticity mechanisms in preterm children with PVL. Furthermore, review articles showed that mechanisms of neuroplasticity after stroke and CP consist of axonal and dendrite sprouting, cortical reorganization and neurogenesis (11).

Nowadays, it is widely accepted that new neurons are formed and recruited into specific brain circuits probably in all adult vertebrate species, including humans (12). Distinct areas of the central nervous system (CNS) are responsible for neurogenesis in the adult brain; the subventricular zone (SVZ) Fig (1), the lining of the lateral ventricles and the subgranular zone (SGZ) of the dentate gyrus within the hippocampus. Neural stem cells (NSCs) are the source of new neurons in the adult brain. The NSCs, which are multi potent and self renewing, are regulated by a number of molecules and signaling pathways.

Certain areas in the brain show that adult neurogenesis continues throughout lifetime. Endogenous progenitor cells located in the subventricular zone enhance this neurogenesis by growth factors or neurotrophic factors, putting forward for consideration that this strategy has the potential to treat the damaged brain. Neural stem cells, neurogenesis, young neuron migration, their differentiation and death have been studied in the subventricular zone-olfactory bulb (SVZ-OB) system which has turned into an appealing experimental model. Furthermore, basic questions are mentioned regarding proliferation and migration of the nerve cells.

Materials & Methods

We searched Medline, CINAHL, Pub Med, ISI Web of Science and Google Scholar from 1995 to February 2011 using a combination of terms such as “Neurogenesis”[Mesh], Stem cell therapy, origin, proliferation dynamics, and migration, brain damage, “Cerebral Palsy”[Mesh], children. The database search identified 51 articles which were retrieved for evaluation. Finally, the results of 48 investigations,

including original and review articles were categorized based on origins, proliferation, and migration of new neurons, and neurogenesis in CP.

Literature Review

There are epidemiological, clinical and review studies about the origins, proliferation, and migration of new neurons, and neurogenesis in children with CP.

Origins

The ceaseless process of producing new neurons in the adult SVZ mentions the possibility of neural stem cell existence within the germinal layer. There are a minimum of four types of cells in the SVZ-ependymal region. These cells have different morphologies, ultrastructures and molecular markers (13) (Fig. 1). Young migratory neurons (type A cells) make up chains which are enclosed in a sheath by astrocytes (type B cells). Type C cells which are more spherical and highly proliferative precursors are gathered together to create migrating A cell chains. Type E cells are ependymal cells that remarkably separate the SVZ from the ventricle cavity. B cells join with E cells intimately leading to occasional contact with the ventricle lumen.

Subventricular zone- astrocytes from the cortex, cerebellum, and spinal cord can also act as stem cells (14) and the primary precursors for new granule neurons (15).

Radial astrocytes (type-B cells) are the primary progenitors in SGZ. They asymmetrically divide in order to produce type-D cells (15). After four stages of maturation, these intermediate progenitors (D1, D2, D2h and D3) finally differentiate into granular neurons (15, 16). Some authors called these SGZ precursors as neuronal progenitors instead of NSCs. The function of these newly generated neurons appears to play a fundamental role in the memory process, learning and depression.

Proliferation

These cells can be proliferated by the administration of growth factors such as epidermal growth factor (EGF) (17), fibroblast growth factor (FGF) (18,19) or their combination (20) into the ventricle after focal

cerebral ischemia. Additionally, newborn neurons are mobilized to migrate not only toward the olfactory bulb (OB) along the rostral migratory stream, but the subependymal overexpression of brain-derived neurotrophic factor (BDNF) also causes neuronal progenitor cells to recruit into a non-neurogenic neostriatum (21,22). The immune system targets neurogenic niches and exerts a considerable effect on proliferation, survival, differentiation and migration of NSCs (23).

Migration

In adult mice, there are millimeters of distance between SVZ and the OB. Doetsch and Alvarez-Buylla reported that "A cells, the SVZ neuroblasts, originate throughout most of the lateral wall of the lateral ventricle and traverse a complex network of interconnected paths before joining the rostral migratory stream (RMS)" (24). It has also been suggested that similar migrations such as migration of SVZ derived cells which is remarkably long may happen in the infant human brain which is large in contrast to mice brain. A study also suggested a migration from the SVZ to neocortex (25, 26), but this migration has not been demonstrated.

A cells in adult rodent SVZ and RMS form chains by moving along each other (Fig. 2) (27). A cells with a long structure and a noticeable leading process are covered at the end by a growth cone (28,29). Neuroblasts move incrementally in chains reconstituted *in vitro* at mean speeds of 120 μ m/hr (28). The cellular construction that causes these young neurons to move at such high speeds is not known. Microtubule polymerization and depolymerization may probably have a significant function in both the exploratory behavior and net translocation happening during a step. Doublecortin, a protein related to microtubule has an important role in neuronal migration.

A cells have an outstanding growth cone and a principal mechanism (28) mentioning that these cells may use motion processes also used by growing axons. A process emerges from the neuron cell body. The forward end of the process expands to form a growth cone that samples the environment, contacting other cells and chemical cues. When the growth cone contacts its target cell, synaptic vesicles soon form and

microtubules that formerly ended at the apex of the growth cone project to the presynaptic membrane.

Besides, during migration and types of signal trigger, factors secreted by astrocytes appear to enhance the migration of SVZ neuroblasts (30). These astrocytes probably play important roles such as enhancement of migration. The glial cells surrounding the chains may also help A cells to survive and/or may provide directional information.

Neurogenesis in CP

Migration of thousands of young neurons (A cells) into the OB happens daily, but only some of them accomplish completely. The characteristics of neural stem cells include the ability to self-renew, differentiate into most types of neurons and glial cells and populate developing and degenerating regions of CNS (31).

Lesions occurring in the perinatal period cause severe impairment of skilled movement learning and also secondary disruption in the development of alpha motor neurons and their afferent segmental reflex control. Because no explicit treatment has been mentioned for CP due to neonatal hypoxic-ischemia (HI), recent research shows that using growth factors can have beneficial effects on ischemic brain injury (32-34).

Neural stem cells and progenitors of the subventricular zone (SVZ) of the adult mammalian brain (35) can be proliferated by the administration of growth factors such as epidermal growth factor (EGF) (17), fibroblast growth factor (FGF) (18,19) or their combination (20,36) into the ventricle after focal cerebral ischemia. Likewise, adult neurogenesis persists in specific brain regions throughout lifetime and can be enhanced from endogenous progenitor cells residing in the SVZ by growth factors or neurotrophic factors, suggesting that this strategy will be able to treat the damaged brain. In other words, if the proliferation and differentiation of newly generated neurons can be directed toward specific functional brain regions, it may be possible to use the recovery of specific functions in the treatment of incurable neurological diseases.

Therefore, induction of striatal neurogenesis by the intraventricular administration of brain derived neurotrophic factor (BDNF) and EGF promoted

functional recovery in an adult animal model of neonatal HI brain injury and this treatment may offer a promising strategy for the restoration of motor function for adults with CP (37).

Besides, stimulation from the external environment could affect the plasticity process. The neural basis of experience-dependent processes appeared to involve active formation of new synaptic connections in response to the events providing the information to be stored (38). Many experiences indicated that nervous system optimizes neural connection during critical periods and lack of necessary experience during critical periods in human development of gross motor and cognition provides retardation in these areas (31). Perhaps using growth factors accompanied with rehabilitation therapy such as neurodevelopmental therapy (NDT) is a good option in the process of neurogenesis.

In this regard, transplantation of several types of stem cells including neural stem cells (39), multipotent adult progenitor cells (40,41) and mesenchymal stem

cells (MSCs) may be beneficial in acute injuries of the CNS (42). Daadi et al. (2010) reported that human neural stem cells (hNSCs) engrafted into the ischemic brain enhanced axonal sprouting and the expression of genes involved in neurogenesis, gliogenesis and neurotrophic support modulated microglial response and improved motor function of the animals. It is generally believed that transplanted non-neural cells such as those derived from bone marrow or cord blood exert neurotrophic effects on ischemia-injured tissue and may not survive long term; (43) whereas, neural stem cells are thought to provide cell replacement and neurotrophic support (44,45).

In the injured brain, growth and differentiation factors released from transplanted MSCs enhance the local trophic milieu; subsequently, improving endogenous repair processes (46,47, 48). It has also been suggested that MSCs can differentiate into neurons and oligodendrocytes and thereby contribute to repair the injured brain (42).

Van Velthoven et al. (2010) suggested that two MSC




Fig 1. The anatomy of the neurogenic subventricular zone in the human brain. (A) Coronal view of the adult human brain showing the basal ganglia and lateral ventricles. (B) Schematic drawing depicting the cellular composition and cytoarchitecture of the adult human SVZ, consisting of four layers: Layer I – ependymal cell layer (green), Layer II – hypocellular gap, Layer III – astrocytic ribbon, containing astrocytes and migrating neuroblasts, Layer IV – transitional zone (From Wikipedia, the free encyclopedia)

injections at 3 and 10 days after neonatal HI markedly improved sensorimotor function 4 weeks after the insult and MSC transplantation after neonatal HI decreased gray and white matter loss and enhanced neurogenesis and oligodendrogenesis. In this study, bone marrow from the femur and tibia of mice was cultured in DMEM/15% fetal bovine serum and was infused into the ipsilateral hemisphere (42). In another study on rats with left carotid artery ligation and hypoxic exposure, it was reported that MSC may be a treatment for neonatal HIE (40).

Obviously, today it is confirmed that using erythropoietin (EPO) with granulocyte colony-

stimulating factor (G-CSF) promoted cell migration in MSCs.

In conclusion, after ischemic brain injury various cell types including neurons, glial and endothelial cells are damaged and lose their function. Effective regeneration of brain tissue requires all these cell types to be replenished and combined to form a new functional network. This review showed that most studies have been conducted in the laboratory and on animals, more work is required at the basic level of stem cell biology, in the development of human models, and finally in well-conceived clinical trials.




Fig 2. Organization and lineage in the SVZ. Left, Cross section of the anterior rodent brain indicating the location of the SVZ on the lateral wall of the LV. On the right is the cellular composition and organization of the SVZ. Chains of young neurons (A cells, red) are surrounded by B cells (blue) that have astrocytic characteristics and form tube-like structures. Ependymal (E), Lateral Ventricle (LV)

References

1. Buonomano DV, Merzenich MM. Cortical plasticity: from synapses to maps. *Annu Rev Neurosci* 1998; 21:149-86.
2. Haghgoo H. *Fundamental of neurosciences*. 1st ed. Tehran; USWR Press; 2011.
3. Payne BR, Lomber SG. Reconstructing functional systems after lesions of cerebral cortex. *Nat Rev Neurosci* 2001 Dec;2(12):911-9.
4. Bax M, Goldstein M, Rosenbaum P, Leviton A, Paneth N, Dan B, et al. Proposed definition and classification of cerebral palsy. *Dev Med Child Neurol* 2005 Apr;47(8):571-6.
5. Joghataei M, Kazem M. Barresi sathe niazhaie jamee

- be khadamate behzisti colle keshvar [persian]. Tehran: University of. Social Welfare and Rehabilitation Sciences; 1990.
6. Johnson A. Prevalence and characteristics of children with cerebral palsy in Europe. *Dev Med Child Neurol* 2002 Sep;44(9):633-40.
 7. Yeargin-Allsopp M, Van Naarden Braun K, Doernberg NS, Benedict RE, Kirby RS, Durkin MS. Prevalence of cerebral palsy in 8-year-old children in three areas of the United States in 2002: a multisite collaboration. *Pediatrics* 2008 Mar;121(3):547-54.
 8. Uvebrant P. Hemiplegic cerebral palsy. Aetiology and outcome. *Acta Paediatr Scand Suppl* 1988;345:1-100.
 9. Marin-Padilla M. Developmental neuropathology and impact of perinatal brain damage. II: white matter lesions of the neocortex. *J Neuropathol Exp Neurol* 1997 Mar;56(3):219-35.
 10. Tzarouchi LC, Astrakas LG, Zikou A, Xydis V, Kosta P, Andronikou S, et al. Periventricular leukomalacia in preterm children: assessment of grey and white matter and cerebrospinal fluid changes by MRI. *Pediatr Radiol* 2009 Dec;39(12):1327-32.
 11. Dehghan L, Dalvand H. Neuroplasticity after stroke [persian]. *Modern rehabilitation* 2008;1(4).
 12. Sanai N, Tramontin AD, Quiñones-Hinojosa A, Barbaro NM, Gupta N, Kunwar S, et al. Unique astrocyte ribbon in adult human brain contains neural stem cells but lacks chain migration. *Nature* 2004 Feb;427(6976):740-4.
 13. Doetsch F, Garcia-Verdugo JM, Alvarez-Buylla A. Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain. *J Neurosci* 1997 Jul 1;17(13):5046-61.
 14. Laywell ED, Rakic P, Kukekov VG, Holland EC, Steindler DA. Identification of a multipotent astrocytic stem cell in the immature and adult mouse brain. *Proc Natl Acad Sci U S A* 2000 Dec 5;97(25):13883-8.
 15. Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Buylla A. Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 2001 Sep 15;21(18):7153-60.
 16. Seri B, Garcia-Verdugo JM, Collado-Morente L, McEwen BS, Alvarez-Buylla A. Cell types, lineage, and architecture of the germinal zone in the adult dentate gyrus. *J Comp Neurol* 2004 Oct 25;478(4):359-78.
 17. Sugiura S, Kitagawa K, Tanaka S, Todo K, Omura-Matsuoka E, Sasaki T, et al. Adenovirus-mediated gene transfer of heparin-binding epidermal growth factor-like growth factor enhances neurogenesis and angiogenesis after focal cerebral ischemia in rats. *Stroke* 2005 Apr;36(4):859-64.
 18. Ellsworth JL, Garcia R, Yu J, Kindy MS. Fibroblast growth factor-18 reduced infarct volumes and behavioral deficits after transient occlusion of the middle cerebral artery in rats. *Stroke* 2003 Jun;34(6):1507-12.
 19. Wada K, Sugimori H, Bhide PG, Moskowitz MA, Finklestein SP. Effect of basic fibroblast growth factor treatment on brain progenitor cells after permanent focal ischemia in rats. *Stroke* 2003 Nov;34(11):2722-8.
 20. Baldauf K, Reymann KG. Influence of EGF/bFGF treatment on proliferation, early neurogenesis and infarct volume after transient focal ischemia. *Brain Res* 2005 Sep 21;1056(2):158-67.
 21. Chmielnicki E, Benraiss A, Economides AN, Goldman SA. Adenovirally expressed noggin and brain-derived neurotrophic factor cooperate to induce new medium spiny neurons from resident progenitor cells in the adult striatal ventricular zone. *J Neurosci* 2004 Mar 3;24(9):2133-42.
 22. Cho SR, Benraiss A, Chmielnicki E, Samdani A, Economides A, Goldman SA. Induction of neostriatal neurogenesis slows disease progression in a transgenic murine model of Huntington disease. *J Clin Invest* 2007 Oct;117(10):2889-902.
 23. Gonzalez-Perez O, Jauregui-Huerta F, Galvez-Contreras AY. Immune system modulates the function of adult neural stem cells. *Curr Immunol Rev* 2010 Aug 1;6(3):167-73.
 24. Doetsch F, Alvarez-Buylla A. Network of tangential pathways for neuronal migration in adult mammalian brain. *Proc Natl Acad Sci* 1996 Dec 10;93(25):14895-900.
 25. Weickert CS, Webster MJ, Colvin SM, Herman

- MM, Hyde TM, Weinberger DR, et al. Localization of epidermal growth factor receptors and putative neuroblasts in human subependymal zone. *J Comp Neurol* 2000 Jul 31;423(3):359-72.
26. Magavi SS, Leavitt BR, Macklis JD. Induction of neurogenesis in the neocortex of adult mice. *Nature* 2000 Jun 22;405(6789):951-5.
27. Lois C, Garcia-Verdugo JM, Alvarez-Buylla A. Chain migration of neuronal precursors. *Science*. 1996 Feb 16;271(5251):978-81.
28. Wichterle H, Garcia-Verdugo JM, Alvarez-Buylla A. Direct evidence for homotypic, glia-independent neuronal migration. *Neuron* 1997 May;18(5):779-91.
29. Lois C, Alvarez-Buylla A. Long-distance neuronal migration in the adult mammalian brain. *Science* 1994 May 20;264(5162):1145-8.
30. Mason HA, Ito S, Corfas G. Extracellular signals that regulate the tangential migration of olfactory bulb neuronal precursors: inducers, inhibitors, and repellents. *J Neurosci* 2001 Oct 1;21(19):7654-63.
31. Lundy-Ekman L. *Neuroscience: fundamentals for rehabilitation*. 3rd ed. Philadelphia: WB Saunders; 2007.
32. Mizuno K, Hida H, Masuda T, Nishino H, Togari H. Pretreatment with low doses of erythropoietin ameliorates brain damage in periventricular leukomalacia by targeting late oligodendrocyte progenitors: a rat model. *Neonatology* 2008;94(4):255-66.
33. Lin S, Fan LW, Pang Y, Rhodes PG, Mitchell HJ, Cai Z. IGF-1 protects oligodendrocyte progenitor cells and improves neurological functions following cerebral hypoxia-ischemia in the neonatal rat. *Brain Res* 2005 Nov 23;1063(1):15-26.
34. Gonzalez FF, Abel R, Almlı CR, Mu D, Wendland M, Ferriero DM. Erythropoietin sustains cognitive function and brain volume after neonatal stroke. *Dev Neurosci* 2009;31(5):403-11.
35. Goldman SA. Adult neurogenesis: from canaries to the clinic. *J Neurobiol* 1998 Aug;36(2):267-86.
36. Tureyen K, Vemuganti R, Bowen KK, Sailor KA, Dempsey RJ. EGF and FGF-2 infusion increases post-ischemic neural progenitor cell proliferation in the adult rat brain. *Neurosurgery* 2005 Dec;57(6):1254-63; discussion -63.
37. Im SH, Yu JH, Park ES, Lee JE, Kim HO, Park KI, et al. Induction of striatal neurogenesis enhances functional recovery in an adult animal model of neonatal hypoxic-ischemic brain injury. *Neuroscience* 2010 Aug 11;169(1):259-68.
38. Greenough WT, Black JE, Wallace CS. Experience and brain development. *Child Dev* 1987 Jun;58(3):539-59.
39. Daadi MM, Davis AS, Arac A, Li Z, Maag AL, Bhatnagar R, et al. Human neural stem cell grafts modify microglial response and enhance axonal sprouting in neonatal hypoxic-ischemic brain injury. *Stroke* 2010 Mar;41(3):516-23.
40. Lee JA, Kim BI, Jo CH, Choi CW, Kim EK, Kim HS, et al. Mesenchymal stem-cell transplantation for hypoxic-ischemic brain injury in neonatal rat model. *Pediatr Res* 2010 Jan;67(1):42-6.
41. Yasuhara T, Matsukawa N, Yu G, Xu L, Mays RW, Kovach J, et al. Behavioral and histological characterization of intrahippocampal grafts of human bone marrow-derived multipotent progenitor cells in neonatal rats with hypoxic-ischemic injury. *Cell Transplant* 2006;15(3):231-8.
42. van Velthoven CT, Kavelaars A, van Bel F, Heijnen CJ. Repeated mesenchymal stem cell treatment after neonatal hypoxia-ischemia has distinct effects on formation and maturation of new neurons and oligodendrocytes leading to restoration of damage, corticospinal motor tract activity, and sensorimotor function. *J Neurosci* 2010 Jul 14;30(28):9603-11.
43. Chopp M, Li Y, Zhang ZG. Mechanisms underlying improved recovery of neurological function after stroke in the rodent after treatment with neurorestorative cell-based therapies. *Stroke* 2009 Mar;40(3 Suppl):S143-5.
44. Buhemann C, Scholz A, Bernreuther C, Malik CY, Braun H, Schachner M, et al. Neuronal differentiation of transplanted embryonic stem cell-derived precursors in stroke lesions of adult rats. *Brain* 2006 Dec;129(Pt 12):3238-48.
45. Daadi MM, Lee SH, Arac A, Grueter BA, Bhatnagar R, Maag AL, et al. Functional engraftment of the medial

- ganglionic eminence cells in experimental stroke model. *Cell Transplant* 2009;18(7):815-26.
46. van Velthoven CT, Kavelaars A, van Bel F, Heijnen CJ. Regeneration of the ischemic brain by engineered stem cells: fuelling endogenous repair processes. *Brain Res Rev* 2009 Jun;61(1):1-13.
47. Qu R, Li Y, Gao Q, Shen L, Zhang J, Liu Z, et al. Neurotrophic and growth factor gene expression profiling of mouse bone marrow stromal cells induced by ischemic brain extracts. *Neuropathology* 2007 Aug;27(4):355-63.
48. Li Y, Chopp M. Marrow stromal cell transplantation in stroke and traumatic brain injury. *Neuroscience Letters* 2009; 456 (3): 120-123.