

Letter to Editor

The Role of Thiazide Diuretics in Treating Idiopathic Hypercalciuria Complications in Children



Dou'aa Albelal¹ , Abolfazl Gilani² , Parsa Yousefchajjan³ , Roham Sarmadian^{4*} 

1. Department of Pediatrics, Faculty of Medicine, Hama University, Hama, Syrian Arab Republic.

2. Department of Pediatric Surgery, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

3. Department of Pediatrics, School of Medicine, Arak University of Medical Sciences, Arak, Iran.

4. Infectious Diseases Research Center (IDRC), Arak University of Medical Sciences, Arak, Iran.



Citation Albelal D, Gilani A, Yousefchajjan P, Sarmadian R. The role of Thiazide Diuretics in Treating Idiopathic Hypercalciuria Complications in Children. Journal of Pediatric Nephrology. 2023; 11(2):114-115. <https://doi.org/10.22037/jpn.v12i2.42592>

 <https://doi.org/10.22037/jpn.v12i2.42592>

Dear Editor,

Idiopathic hypercalciuria (IHC) is a prevalent disorder during childhood and a significant contributor to the formation of renal stones and urinary tract infections (UTIs). Thiazide diuretics, such as hydrochlorothiazide (HCT), are widely accessible, affordable, and easy to administer, with minimal side effects. These characteristics make them acceptable pharmacological options for treating hypercalciuria. The use of thiazides enhances calcium reabsorption from distal renal tubules, reducing hypercalciuria, and potentially preventing associated complications.

Numerous studies have confirmed the beneficial role of HCT in addressing various complications associated with IHC in pediatric patients.

HCT was administered to 59 children with IHC and recurrent UTI, resulting in no UTI recurrence for 95% of these patients [1]. Another single-blind randomized clinical trial involving 100 girls aged 1 to 12 years with IHC and at least two UTIs within a year demonstrated that all patients treated with HCT became normocalciuric. However, treating hypercalciuria did not have a significant impact on UTI prevention [2]. Consequently, further exploration in this area is warranted.

Preliminary research indicates that the long-term use of thiazide diuretics is correlated with decreased recurrent

kidney stone incidence. Nevertheless, due to limited evidence and the economic burden of chronic medication, these drugs have not been advised for preventing recurrent kidney stones [3].

In children with IHC, HCT has been shown to positively affect bone mineral density [4]. Thiazides have positive effects on osteoblastic cell proliferation and activity. Thiazide therapy may improve bone formation and mineralization by inhibiting osteocalcin expression by osteoblasts [5]. In contrast, García-Nieto et al. determined that children with IHC and osteopenia who were given thiazides for 2.4 years had no improvement in bone mineral density (z-BMD) compared to controls [6]. The results may have been confounded by growth during the study period.

Another aspect to consider in the relationship between IHC and HCT is the potential effect of drugs on nocturnal enuresis. Children with IHC are commonly prescribed HCT to treat nocturnal enuresis, and our study confirmed that a single daily dose of HCT (1 mg/kg/day) was a safe and efficient treatment option [7].

As a result, HCT seems to be an effective and safe medication for treating IHC-related complications in children. However, there are still controversies regarding the effect of HCT on complications, such as UTI and osteopenia in IHC, and additional research is required to gain a more comprehensive understanding of the drug's efficacy and safety in larger populations.

Corresponding Author:

Roham Sarmadian

Address: Infectious Diseases Research Center (IDRC),
Arak University of Medical Sciences, Arak, Iran.

E-mail: rsarmadian@yahoo.com

References

- [1] López MM, Castillo LA, Chávez JB, Ramones C. Hypercalciuria and recurrent urinary tract infection in Venezuelan children. *Pediatr Nephrol.* 1999; 13(5):433-7. [DOI:10.1007/s004670050635] [PMID]
- [2] Yousefi P, Firouzifar M, Cyrus A. Does hydrochlorothiazide prevent recurrent urinary tract infection in children with idiopathic hypercalciuria? *J Pediatr Urol.* 2013; 9(6 Pt A):775-8. [DOI:10.1016/j.jpuro.2012.09.008] [PMID]
- [3] Li DF, Gao YL, Liu HC, Huang XC, Zhu RF, Zhu CT. Use of thiazide diuretics for the prevention of recurrent kidney calculi: A systematic review and meta-analysis. *J Transl Med.* 2020; 18(1):106. [DOI:10.1186/s12967-020-02270-7] [PMID]
- [4] Penido M, Tavares MS. Should pediatric idiopathic hypercalciuria be treated with hypocalciuric agents? *World J Nephrol.* 2021; 10(4):47-58. [DOI:10.5527/wjn.v10.i4.47] [PMID]
- [5] Ryan LE, Ing SW. Idiopathic hypercalciuria and bone health. *Curr Osteoporos Rep.* 2012; 10(4):286-95. [DOI:10.1007/s11914-012-0120-5] [PMID]
- [6] García-Nieto V, Monge-Zamorano M, González-García M, Luis-Yanes MI. Effect of thiazides on bone mineral density in children with idiopathic hypercalciuria. *Pediatr Nephrol.* 2012; 27(2):261-8. [DOI:10.1007/s00467-011-1987-6] [PMID]
- [7] Yousefichaijan P, Sharafkhan M, Cyrus A, Rafeie M. Therapeutic efficacy of hydrochlorothiazide in primary monosymptomatic nocturnal enuresis in boys with idiopathic hypercalciuria. *Nephrourol Mon.* 2015; 7(5):e29127. [DOI:10.5812/numonthly.29127] [PMID] [PMCID]