

Mesothelioma in Situ of the Spermatic Cord Arising from a Patent Processus Vaginalis: A Case Report

Yasuyuki Kobayashi^{1*}, Yumiko Yasuhara², Hiroki Arai¹, Masahito Honda¹, Masataka Hiramatsu³, Sho Goya³

Mesothelioma is an aggressive tumor originating from mesothelial cells. Mesothelioma of the spermatic cord is a very rare disease, and the most common presentation of this disease is that of aggressive mesothelioma with no description of mesothelioma in situ. We report an extremely rare case of mesothelioma in situ of the spermatic cord arising from a patent processus vaginalis. To our best knowledge, this is the first report of this finding. The identification of a patent processus vaginalis and investigation of single-layered atypical mesothelial cells led to the final diagnosis.

Keywords: atypical mesothelial cells; mesothelioma in situ; radical orchiectomy; spermatic cord tumor

INTRODUCTION

Mesothelioma is an aggressive tumor originating from mesothelial cells⁽¹⁾. Mesothelioma of the spermatic cord is a rare disease, with only 9 cases reported in detail in the English literature and none describing mesothelioma in situ⁽²⁻¹⁰⁾. We report an extremely rare case of mesothelioma in situ of the spermatic cord arising from a patent processus vaginalis.

CASE REPORT

An 82-year-old man complaining of a gradually enlarging swollen mass in his right inguinal area for the past three months was referred to our hospital. The right spermatic cord mass was hard, smooth, and immobile. Chest and abdominal computed tomography showed right spermatic cord swelling and no other significant findings (**Figure 1**). We diagnosed the patient as having right spermatic cord tumor, for which he underwent right radical orchiectomy. A spermatic cord tumor was found adhering to the surrounding tissue, and no obvious residual tumor existed intraoperatively. Macroscopically, the spermatic cord was enlarged at 7.0 × 4.0 cm and had a smooth and capsulated surface. The cut surface was creamy white, solid, thickened and poorly circumscribed (**Figures 2A, B**). Microscopically, the tumor was a mixture of tubulopapillary, trabecular, and solid structures (**Figures 2C, D**). Immunohistochemically, the tumor cells were positive for calretinin, Wilms' tumor 1, podoplanin, glucose transporter type 1 (GLUT-1) and epithelial membrane antigen (EMA), negative for carcinoembryonic antigen, MOC-31 and Ber-EP4, and showed loss of BRCA1-associated protein (BAP1). We diagnosed the spermatic cord tumor as epithelial mesothelioma. Interestingly, a patent processus vaginalis was found lined by a single layer of mesothelial cells from the proximal end to the tunica vaginalis (**Figure 3A**). A single layer of atypical mesothelial cells with enlarged nuclei lined the ipsilateral side of the mesothelioma and flattened normal mesothelial cells lined the cavity on the opposite




Figure 1. Abdominal CT showing the right spermatic cord swelling.

Departments of Urology, 2Diagnostic Pathology, and 3Respiratory Medicine, Kinki Central Hospital of Mutual Aid Association of Public School Teachers, Itami, Hyogo, Japan.

*Correspondence: Department of Urology, Kinki Central Hospital of Mutual Aid Association of Public School Teachers, 3-1 Kurumazuka, Itami, Hyogo 664-8533, Japan

Phone: +81-72-781-3712; Fax: +81-72-779-1567. E-mail: ya_su_koba@yahoo.co.jp.

Received June 2019 & Accepted December 2019




Figure 2. (A) Macroscopic findings of the spermatic cord. The spermatic cord was enlarged (arrowheads). (B) The cut surface showed a solid and thickened spermatic cord (arrowheads) and no involvement of the testis and tunica vaginalis (arrows). (C) Epithelioid mesothelioma with a papillary pattern of growth. (D) Epithelioid mesothelioma with trabecular and solid patterns of growth. (C-D): Hematoxylin and eosin stain.

side (Figure 3B). Immunohistochemically, the atypical mesothelial cells were positive for GLUT-1 and EMA, negative for desmin, and showed loss of BAP1, whereas the normal mesothelial cells showed an opposite pattern (Figures 3C-D), indicating that the single-layered atypical mesothelial cells were mesothelioma in situ. Finally, we diagnosed mesothelioma in situ of the spermatic cord arising from a patent processus vaginalis. A sec-

ond surgery and adjuvant therapy were not performed because there was no obvious tumor elsewhere. He was alive without signs of disease at 2 years after surgery.

DISCUSSION

Mesothelioma of the spermatic cord is a rare entity; only 9 cases have been reported so far in the English




Figure 3. (A) Cross-sections of the specimen of the right spermatic cord. (a) Mesothelioma of the spermatic cord (red line), (b) patent processus vaginalis (yellow line) and (c) vas deferens (green point). Patent processus vaginalis stained with hematoxylin and eosin (B), desmin (C) and BAP-1 (D). (B-D) Mesothelioma is present on the right side of the images. Cells on the right lining are mesothelioma in situ (arrowhead). Cells on the left lining are normal mesothelial cells (arrow). ※ = patent processus vaginalis.

Table 1. Clinical and pathologic features of primary mesothelioma.

Case	Age (Y)	Asbestos exposure	Side	Symptom	Tumor metastasis	Operation	Histologic type	Postoperative recurrence	Follow-up	Ref
1	63	NA	Lt	Mass in the inguinal region	NA	NA	Biphasic	NA	NA	2
2	40	NA	Lt	High near groin	External iliac lymph node	RO, RALD	NA	No	NED 18 Y	3
3	57	NA	Rt	Inguinal hernia	No	SR	Epithelioid	Local at 12 M Periaortic, iliac lymph nodes at 34 M	DOD 42 M	4
4	46	No	Rt	Mass in the inguinal region	No	RA	Biphasic	NA	DOD 8 M	5
5	52	NA	Rt	Scrotal enlargement	No	RA	Epithelioid	Local at 9 M Iliac lymph nodes at 30 M	AWD 30 M	6
6	65	Yes	Lt	Mass in the inguinal region	No	RA	Biphasic	Local, peritoneal at 3 M	DOD 6 M	7
7	45	No	Lt	Mass in the inguinal region	No	SR	Epithelioid	No	NED 6 M	8
8	80	Yes	Rt	Mass in the inguinal region	No	RA	Biphasic	No	NED 12 M	9
9	45	No	Rt	Swelling in the inguinal region	No	RA	Biphasic	NA	NA	10
Our case	82	No	Rt	Swelling in the inguinal region	No	RA	Epithelioid	No	NED 2 Y	

Abbreviations: NA, not available; Lt, left; Rt, right; RO, radical orchiectomy; RALD, retroperitoneal and para-aortic lymph node dissection; SR, spermatic cord resection; NED, no evidence of disease; DOD, dead of disease; AWD, alive with disease; Y, years; M, months; Ref, reference.

literature (Table 1)⁽²⁻¹⁰⁾. The patients most commonly presented with an inguinal mass, and all 9 patients underwent surgery. Histological findings showed no descriptions of mesothelioma in situ. Mesothelioma arising from the peritoneum may have invaded the spermatic cord in Cases 4 and 6. Our patient being alive without signs of disease at 2 years after surgery excluded the possibility of a peritoneal origin. BAP1 expression is detected by immunohistochemistry, which is useful for distinguishing benign and malignant mesothelial proliferations⁽¹¹⁾. BAP1 expression is detected in the nuclei of benign mesothelial cells, whereas BAP1 loss is detected in the nuclei of mesothelioma cells⁽¹²⁾. Recently, Churg et al. reported two cases of mesothelioma in situ. They proposed that mesothelioma in situ be defined as well-confirmed in situ lesions and be related to genomic events such as BAP1 loss⁽¹³⁾. The single-layered atypical mesothelial cells in our patient were an in situ lesion and showed loss of BAP1. Thus, they were diagnosed as mesothelioma in situ of the spermatic cord arising from a patent processus vaginalis.

REFERENCES

- Gemba K, Fujimoto N, Kato K, et al. National survey of malignant mesothelioma and asbestos exposure in Japan. *Cancer Sci.* 2012; 103:483-90.
- Kozłowski H, Zoltowska A. Mesothelioma of spermatic cord. *Neoplasma.* 1968; 15:97-100.
- Arlen M, Grabstald H, Whitmore WF Jr. Malignant tumors of the spermatic cord. *Cancer.* 1969; 23:525-32.
- Pizzolato P, Lamberty J. Mesothelioma of spermatic cord: electron microscopic and histochemical characteristics of its mucopolysaccharides. *Urology.* 1976; 8:403-8.
- Leiber C, Katzenwadel A, Popken G, Kersten A, Schultze-Seemann W. Tumour of the spermatic cord: an unusual primary manifestation of an epithelial mesothelioma of the peritoneum with patent processus vaginalis. *BJU Int.* 2000; 86:142-3.
- Torbati PM, Parvin M, Ziaee SA. Malignant mesothelioma of the spermatic cord: case report and review of the literature. *Urol J.* 2005; 2:115-7.
- Park YJ, Kong HJ, Jang HC, Shin HS, Oh HK, Park JS. Malignant mesothelioma of the spermatic cord. *Korean J Urol.* 2011; 52:225-9.
- Meng X, Guzzo TJ, Bing Z. Malignant mesotheliomas in spermatic cords: reports of two cases and a brief review of literature. *Rare Tumors.* 2013; 26:5:e4.
- D'Antonio A, Mastella F, Colucci A, Silvestre G. Malignant mesothelioma of spermatic cord in an elderly man with a history of asbestos exposure. *Urology.* 2016; 87:e1-3.
- Ahmed Z, Singh S, Mangal A, Mittal A. Primary malignant mesothelioma of the spermatic cord. *BMJ Case Rep.* 2016; pii: bcr2016214602.
- Kinoshita Y, Hida T, Hamasaki M, et al. A combination of MTAP and BAP1 immunohistochemistry in pleural effusion cytology for the diagnosis of mesothelioma. *Cancer Cytopathol.* 2018; 126:54-63.
- Wu D, Hiroshima K, Yusa T, et al. Usefulness of p16/CDKN2A fluorescence in situ hybridization and BAP1 immunohistochemistry for the diagnosis of biphasic mesothelioma. *Ann Diagn Pathol.* 2017; 26:31-7.
- Churg A, Hwang H, Tan L, et al. Malignant mesothelioma in situ. *Histopathology.* 2018; 72:1033-8.