

Running head: ROBOTIC EXCISION OF THE VAGINA IN A PEDIATRIC PATIENT

Robotic Excision of the Vagina in a 46 XX DSD Male Patient. First Pediatric Report

Authors:

Giovanni Torino MD, Pediatric Urology Unit, "Santobono-Pausilipon" Children's Hospital, Naples (Italy)

Ottavio Adorisio MD, Department of Pediatric Surgery, Pediatric Surgery Unit, Bambino Gesù Children's Hospital, Research Institute, Palidoro, Rome, Italy

Giovanni Cobellis MD, Pediatric Surgery and Urology Unit, "G. Salesi" Children's Hospital, Marche Polytechnic University, Ancona, Italy

Francesca Mariscoli MD, Pediatric Surgery and Urology Unit, "G. Salesi" Children's Hospital, Marche Polytechnic University, Ancona, Italy

Antonio Zaccara MD, Department of Pediatric Surgery, Pediatric Urology and Urodynamics Unit, Bambino Gesù Children's Hospital, Research Institute, Palidoro, Rome, Italy

Key words: Robotic surgery; DSD; UTI; pelvic surgery; minimally invasive surgery.

Abstract

The Disorders of Sex Differentiation (DSD) represent a wide range of congenital anomalies of the genitalia. Surgical treatment of these cases may become a challenge. We present a case of a 16-year-old boy with 46 XX DSD, SRY negative, presented with persistent dribbling incontinence, recurrent UTI and perineal pain. Past medical history included right orchiectomy, laparoscopic excision of uterus, fallopian tubes and left streak gonad at another institution at the age of 2 years. The native vagina was left in place. VCUG confirmed the presence of the residual vagina (8 cm in maximum length), connected with the bulbar urethra. Robotic-assisted laparoscopy of the vagina was performed with satisfying short and long-term results.

Introduction

Robotic removal of the vagina in a pediatric male patient has not yet been reported to date. We report a case of a 16-year-old boy affected by 46 XX DSD with a residual vagina removed using the robotic approach.

Case Report

A 16-year-old boy with 46 XX DSD, SRY negative, presented with persistent dribbling incontinence, recurrent febrile UTI and perineal pain. He was raised as a male by his parents. His past medical history included: right inguinal orchiectomy for ovotestes, laparoscopic excision of the uterus, fallopian tubes and intra-abdominal left streak gonad at another institution at the age of 2 years (the native vagina was left in place), two-stage repair of scrotal hypospadias when he was 3-year-old, bilateral mastectomy at the age of 13 years. Because of the persistence of recurrent febrile UTI and perineal pain, a voiding cystourethrogram (VCUG) was performed, confirming the presence of a residual vagina (8 cm maximum

length) (Fig. 1). The decision to perform a vaginectomy was taken. The patient was placed in a combined lithotomy/supine position to perform simultaneously the robotic surgery and the vaginoscopy. The preliminary cystoscopy showed a normal urethra, without stenosis, with the vagina opening at the level of the bulbar urethra. The 12-mm optical port was inserted via the umbilicus, and other two robotic 8-mm working ports were placed on the para-rectal lines and at the level of the transverse umbilical line. The bladder was suspended to the anterior abdominal. The light of the cystoscope, into the vagina, was used as guidance to separate the vagina from both the surrounding tissues (Fig 2A-B). A dissection was performed between the vagina and the surrounding tissues. The suspension of the vagina, the good magnification ensured by the robotic approach, and the Trendelenburg position allowed us to avoid injuries of the sacral plexus. The vagina was transected as close as possible to the urethra leaving in place a little sidewall of the vagina near the urethra. This opening was closed with an absorbable stitch, avoiding urethral strictures (Video 1). The specimen was then removed from the optical port. The pathologic examination confirmed the removal of the vagina. The operative time was 240 minutes.

The post-operative period was uneventful. The patient was discharged 7 days after the procedure. The Foley, previously inserted, was removed on the 20th post-operative day. A post-operative VCUG showed neither significant remnants at the level of the bulbar urethra or leakage with normal bladder emptying (Fig. 3). After a two-year follow-up, the patient shows a complete resolution of the symptoms.

Discussion

Surgical removal of a residual vagina may be a real challenge in the pediatric population due to the proximity to the pelvic nerves, rectum, ureters, bladder,

rectal sphincters and, because of the narrow surgical space (1,2,3). There is a gaining consensus about robotic surgery feasibility and efficacy in the pediatric population due to an increasing number of appreciable surgical results reported in the literature in a wide range of congenital anomalies (1). In the case of pelvic anomalies, open surgery needs often of a combined approach with two incisions at the level of the abdomen and perineum respectively. The laparoscopic approach, despite shorter hospitalization and good control of post-operative pain, rarely allow to safely reach the perineum space (4,5). The minimally invasive approach, based on robotic surgery, has been recommended in several reports (4,6,7) because of the clearer view of the deep pelvic structures. Robotic-assisted surgery has been successfully used to obtain a complete removal of a large utricle in a pediatric patient who previously undergone to the laparoscopic procedure (8). In the present case, the previously laparoscopic approach did not allow the removal of the vagina, for this reason, we opted for robotic-assisted laparoscopy in the redo procedure to combine the advantages of laparoscopy with the improved three-dimensional (3D) visualization associated with high instruments dexterity (9). This technique has been previously used in transgender adults (10). To our knowledge, no cases of robot-assisted excision of the residual vagina in a pediatric DSD male patient has been described in the literature to date.

Conclusions

In conclusion, because of the excellent magnification, the high dexterity ensured by robotic instruments, the low risk of iatrogenic lesions, the short hospital stay and the low postoperative pain, robotic-assisted surgery should be considered also in pediatric patients.

Conflict of interest: The Authors declare no conflicts of interest

References

1. Goruppi I, Avolio L, Romano P, Raffaele A, Pelizzo G. Robotic-assisted surgery for excision of an enlarged prostatic utricle. *Int J Surg Case Rep.* 2015;10:94-6.
2. Alqahtani A, Albassam A, Zamakhshary et al. Robot-assisted pediatric surgery: how far can we go? *World J Surg,* 2010;34:975-8.
3. Matthews CA. Applications of robotic surgery in gynecology. *J Womens Health (Larchmt),* 2010;19:863-7.
4. Onal B, Diamond DA, Retik AB, Cendrom M, Nguyen HT. Robot-assisted laparoscopic excision of symptomatic retrovesical cyst in boys and young adults. *J Urol.* 2011;186:2372-2378.
5. Estrada CR, Passerotti CC. Robotic surgery in pediatric urology. *Arch. Esp. Urol.* 2007;60:471-479.
6. Benmohamed B, Chaker K, Obringer L et al. Duplicated collecting system with ectopic prostatic implantation: Therapeutic conservative robot assisted approach, about a case report. *Urol Case Rep.* 2019 Nov 6;30.
7. Lima M, Maffi M, Di Salvo N et al. Robotic removal of Müllerian duct remnants in pediatric patients: our experience and a review of the literature. *Pediatr Med Chir.* 2018 May 30;40.
8. Ferong K, Waterschoot M, Sinatti C et al. Rare and special robotic surgery indications in the pediatric population: ectopic

organs and differences of sexual development. World J Urol. 2019 Aug 22.

9. Benmohamed B, Chaker K, Obringer L et al. Duplicated collecting system with ectopic prostatic implantation: Therapeutic conservative robot assisted approach, about a case report. Urol Case Rep. 2019 Nov 6;30.
10. Cohen OD, Dy GW, Nolan IT, Maffucci F, et al. Robotic Excision of Vaginal Remnant/Urethral Diverticulum for Relief of Urinary Symptoms Following Phalloplasty in Transgender Men. Urology. 2020;136:158-161.

Corresponding author:

Dr. Ottavio Adorisio

Department of Pediatric Surgery

Children's Hospital "Bambino Gesù Children's Hospital, via della torre di

Palidoro 50, Palidoro, Rome, Italy

Legends to figures:

Figure 1: Preoperative VCUG showing the vagina (8 cm), filling up during micturition.

Figure 2 A-B: Separation of the vagina from the surrounding structures using the light of the cystoscope as a guidance.

Figure 3: Post-operative VCUG showing the removal of the vagina with a little stump close to the urethra.



Figure 1: Preoperative VCUG showing the vagina (8 cm), filling up during micturition.




Figure 2 A-B: Separation of the vagina from the surrounding structures using the light of the cystoscope as a guidance.



Figure 3: Post-operative VCUG showing the removal of the vagina with a little stump close to the urethra.

Accepted