

Post Circumcision “Tri-Balanic” Penis

*Urol J. 2011;8:13.
www.uj.unrc.ir*



A 32-year old man presented to the outpatient clinic of our department complaining of intractable pain, considerable congestion, and edema of the glans of the penis. The patient was afebrile, complaining of dribbling and inability to urinate properly.

The patient’s history included a recent (less than one month) operation for the penis circumcision in another institute, while the physical examination revealed glans lobulated and struggled.

Under general anesthesia, the stitches were released and the redundant part of the prepuce was excised as to release the compression of the glans, leaving the glans free and totally uncovered. Following this procedure, the edema and congestion improved.

This case represents a rare case of circumcision repair-induced paraphimosis. Rules to be followed in order to avoid this complication are removal of adequate part of the prepuce, careful detachment of adhesions between the inner surface of the prepuce and superficial surface of the glans,⁽¹⁾ gentle removal of the smegma, extreme pull back of the prepuce behind the corona of the glans,⁽²⁾ and careful ligation of the artery and veins.⁽³⁾

Konstantinos Michalakis, Ioannis Ilias, Charalambos Asvestis

*Elena Venizelou General Hospital, Athens, Greece
E-mail: kostismichalakis@hotmail.com*

REFERENCES

1. Elder JS. Circumcision. *BJU Int.* 2007;99:1553-64.
2. Muula A, Prozesky H, Mataya R, Ikechebelu J. Prevalence of complications of male circumcision in Anglophone Africa: a systematic review. *BMC Urol.* 2007;7:4.
3. Holman J, Stuessi K. Adult circumcision. *Am Fam Physician.* 1999;59:1514-8.